


Innovative switching mechanism improves ultrafast control of microlasers

February 28 2020



Ultrafast control of the quasi-BIC microlasers. (A) Schematic of two-beam pumping experiments. Two beams are spatially detuned with a distance d . Science (2020). DOI: 10.1126/science.aba4597

The all-optical switch is a kind of device that controls light with light, which is the fundamental building block of modern optical communications and information processing. Creating an efficient,

ultrafast, and compact all-optical switch has been recognized as the key step for the developments of next-generation optical and quantum computing. In principle, photons don't interact with one another directly in the low power linear regime, and a cavity is usually needed to resonantly enhance the field of control light and increase the interaction. In early work, the performance of all-optical switches has been improved rapidly by optimizing resonators such as microrings or photonic crystals. For further improvements, the research area reaches the limit—the trade-off between ultralow energy consumption and ultrashort switching time.

"Low energy consumption usually requires a high Q factor of the resonator, whereas the longer lifetime high-Q mode imposes an obstacle for improving switching speed," said Qinghai Song from Harbin Institute of Technology, China. "An alternative approach with plasmonic nanostructure has been recently exploited to break the trade-off. The inserting and propagating loss is as large as 19 dB and additional power consumption is required to amplify the signals."

The lasing actions at the topologically protected bounded states in the continuum has the potential to eventually solve this long-standing challenge. In *Science*, researchers from Harbin Institute of Technology, Australian National University and City University of New York detail their innovation of the switching mechanism at the topologically protected bounded states in the continuum (BICs), which offers an ultrafast transition of microlaser emission from a radially polarized donut beam to linearly polarized lobes and vice versa. The extremely high Q factor of the BICs can dramatically reduce the [laser](#) threshold and eventually break the above trade-off in conventional all-optical switches.

The next step of this research is to integrate cascade-wise several such switchable microlasers with an integrated photonic chip and to perform

optical logic operations. This is the prerequisite for the ultimate goal—optical or quantum computing.

Using symmetry-protected BICs

Conventional ultrafast all-optical switches either utilize the nonlinear refractive index or nonlinear absorption to produce an optical bit. Such techniques require either high excitation fluence to produce wavelength shift or extra loss to modulate the transmission, still limited by the trade-off.

The researchers solve this problem with the optical characteristics of BICs, which was initially proposed by Von Neumann and Wigner in quantum mechanics and has been revisited in optics recently. Although the accidental BICs has attracted most research attention due to its robustness, this research focuses on laser emissions from symmetry protected BICs. Compared with accidental BICs, the latter is extremely sensitive to symmetry-breaking perturbations. The exceptional gain of a lasing system, corresponding to the imaginary part of the refractive index (n''), can be a new and effective parameter to ultrafast control of symmetry and the corresponding symmetry-related far-field laser emissions at the BICs.

To test the concept, the researchers fabricated a square lattice periodic nanostructure in a MAPbBr_3 perovskite film and optically pumped it. Single-mode laser operation was achieved at the symmetry protected BICs. The output laser beam is a donut beam in the vertical direction with a divergence angle of 2θ . The polarization test and self-interference pattern show the emission laser beam is radially polarized with orbital angular momentum (OAM).

"The directional laser emission with OAM is not surprising," said Song. "It has been observed and explained by B. Kante from UC Berkeley. It

relates to the polarization vortex at the BICs and the transverse spin angular momentum induced by the real samples. It can also be realized with circularly polarized optical excitation. Comparably, the BIC microlaser is more intriguing in all-optical switching."

The researchers show that the pumping profile can effectively control the BIC lasers. By partially increasing the optical gain with the second beam, the four-fold rotational symmetry is broken and the BIC laser degrades to a conventional photonic crystal laser. As a result, the donut beam transits to two linearly polarized lobes. Such a transition and its reversal happen in a time of 1-1.5 ps. A complete transition from a donut to two lobes and back to a donut has also been realized within 2-3 ps. Such a switching time is more than an order of magnitude faster than the BIC microlaser lifetime, clearly demonstrating that the limitation of laser lifetime on switching time is broken.

"Such an ultrafast control is attributed to the far-field characteristics of BICs," said Song. "The BICs are formed by destructive interference at the radiation channels. Considering far field radiation, the transition from BIC microlasers to conventional lasers represents a re-distribution of the laser emission instead of a direct switching on/off of the lasing mode."

The laser threshold is about 4.2 mW/cm^2 , giving an energy consumption per bit similar to current all-optical switches. "This is because the quality of our perovskite nanostructures is low and the ultrahigh Q factor of BICs has not been fully utilized," said Song. "Eventually, the threshold can be further reduced by orders of magnitude with the BICs and break all the limitations of all-optical switches."

The demonstrated mechanism is not limited by optical excitation. Electrically-driven BIC microlasers with ultrafast switching are also possible and the cascade-wise on-chip integration of such ultrafast

switchable BIC lasers is also essential for optical and quantum computing. This research is published in *Science* on Feb. 28, 2020.

More information: Can Huang et al. Ultrafast control of vortex microlasers, *Science* (2020). [DOI: 10.1126/science.aba4597](https://doi.org/10.1126/science.aba4597)

Provided by Harbin Institute of Technology

Citation: Innovative switching mechanism improves ultrafast control of microlasers (2020, February 28) retrieved 3 May 2024 from <https://phys.org/news/2020-02-mechanism-ultrafast-microlasers.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|