

# Mediterranean sea urchins are more vulnerable than previously thought

February 7 2020

---



The species *Paracentrotus lividus* is very abundant in the Catalan coasts. Credit: Creu Palacín (University of Barcelona-IRBio)

The sea urchin *Paracentrotus lividus*, an eatable species of great commercial interest found in the Mediterranean and North-East Atlantic, is more vulnerable than so far believed.

This is stated in a study by the University of Barcelona, the Spanish National Research Council (CSIC) and the University of Tromsø (Norway) on the genetic distribution of populations of this species, led by the experts Marta Pascual, from the Faculty of Biology and the Biodiversity Research Institute (IRBio) of the University of Barcelona, and Xavier Turon, CSIC research professor at the Center for Advanced Studies of Blanes (CEAB-CSIC). The study has been recently published in the journal *Diversity and Distributions*.

Using a genetic analysis of the populations of this sea urchin, this work assesses its [population structure](#) and connectivity as a result of its dispersal ability. Scientists analysed about 250 individuals of *Paracentrotus lividus* from eleven marine areas, ranging from the French Atlantic coast to the eastern Mediterranean.

Researcher Xavier Turon (CEAB-CSIC) notes that "adult sea urchin barely move," and that dispersal occurs during the larval stage, since "when they are larvae, they swim in the plankton and are carried away by marine currents. We thought they had a high dispersal capability, since they have a long larval lifespan, which lasts for weeks."

What the study reveals is that "the dispersal is not as wide as we expected," says Turon, "and geographical barriers such as the Strait of Gibraltar limit the connectivity of populations. The exchange of sea urchin larvae between different areas is more limited than what we thought."

Another shocking result is that the selection at a local level is having an impact on the genetic make-up of the different populations of urchins.

"We found different allele composition among populations," note researchers Marta Pascual and Carles Carreras (UB-IRBio). "When we analysed the genetic sequences in which these differences appeared, we saw that they lay in genes that code for processes related to temperature and salinity." There is a strong selection pressure related to temperature in the Mediterranean, which can have important consequences regarding global warming.

The study reveals there is a progressive adaptation to salinity and temperature from west to east, and that the largest genetic differences are found between populations in the Atlantic and the Mediterranean. In the latter, the populations of western Mediterranean areas are also differentiated from those in the eastern Mediterranean.

Despite being the target of an active fishery, this species is very abundant in the Catalan coasts, since the fishing pressure has removed many of its predators (mainly fish). Urchins are herbivorous and eat seaweeds and seagrasses. With no predators, the populations can grow excessively and can eliminate submarine forests by overgrazing.

There are two main species of sea urchins in the littoral area of the Mediterranean: the eatable species *Paracentrotus lividus*, and the black sea urchin *Arbacia lixula*, non-edible. The new study brings new data to consider so as to predict potential future scenarios of how the common sea urchin populations will respond to antagonist pressures of predator removal and fishing, in a context of progressive warming.

**More information:** Carlos Carreras et al, East is East and West is West: Population genomics and hierarchical analyses reveal genetic structure and adaptation footprints in the keystone species *Paracentrotus lividus* (Echinoidea), *Diversity and Distributions* (2019). [DOI: 10.1111/ddi.13016](https://doi.org/10.1111/ddi.13016)

Provided by University of Barcelona

Citation: Mediterranean sea urchins are more vulnerable than previously thought (2020, February 7) retrieved 23 April 2024 from

<https://phys.org/news/2020-02-mediterranean-sea-urchins-vulnerable-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.