

Secondary forests provide deforestation buffer for old-growth primary forests

February 6 2020



Secondary forest conservation, July, 2014 in São Félix do Xingu. Credit: Nelton Cavalcante da Luz & Douglas Rafael Moraes Vidal

Currently, re-growing forests comprise roughly 21% of previously deforested areas in the Brazilian Amazon. However, these forests, referred to as secondary vegetation, have been little studied, despite occupying a total area similar to that of the United Kingdom.

Now, researchers led by the University of Leeds, the Brazilian National Institute for Space Research (INPE), and Brazilian Agricultural Research Corporation (EMBRAPA) have examined 14 years of data on secondary vegetation formation and cutting in the Brazilian Amazon based on the TerraClass Amazon mapping project.

Their study, published in *Nature Sustainability*, has found that [secondary forests](#) account for an increasing proportion of overall forest loss in the Brazilian Amazon, rising from 32% of total deforestation in 2000 to 72% in 2014.

However, while secondary forest cutting has increased in recent years, deforestation of old-growth primary forests in the Brazilian Amazon remained stable. This suggests that secondary forest loss has eased deforestation pressure on primary forests and their irreplaceable biodiversity and carbon storage.

Study lead author Yunxia Wang, from the School of Geography said: "There is an ongoing demand for new pasture and agricultural land in Brazil. Our study shows that this demand has increasingly been met by secondary forests, providing a buffer that has stalled deforestation of primary forests.

"But the strength of this buffer depends on the area of secondary forest available. The limited legal protection means that secondary forest loss is largely unregulated.

"Not only would easing the strain on secondary forests help Brazil meet

climate change targets, as they accumulate carbon very rapidly, but future [deforestation](#) would likely lead to increased loss of primary forests once easily accessible secondary forests are diminished."

Brazil has committed to restore 120,000 km² of forest land by 2030 as part of its Nationally Determined Contribution for the Paris Agreement. The authors suggest that a cost-effective way to do this would be to allow part of its existing Amazonian secondary [forest](#) area to recover naturally.

Study co-author Dr. David Galbraith, Associate Professor in Earth System Dynamics at Leeds said: "Managing this ecosystem sustainably to maximize the conservation value of these forests, while not intensifying pressure on [primary forests](#), requires an integrated strategy that includes active monitoring of secondary forests in Amazonia and strengthening of their governance in Brazilian law."

More information: Yunxia Wang et al. Upturn in secondary forest clearing buffers primary forest loss in the Brazilian Amazon, *Nature Sustainability* (2020). [DOI: 10.1038/s41893-019-0470-4](https://doi.org/10.1038/s41893-019-0470-4)

Provided by University of Leeds

Citation: Secondary forests provide deforestation buffer for old-growth primary forests (2020, February 6) retrieved 3 August 2024 from <https://phys.org/news/2020-02-secondary-forests-deforestation-buffer-old-growth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.