


Like snowflakes, soot particles are unique, affecting climate modeling

February 27 2020



Credit: Brookhaven National Laboratory

Black carbon particles—more commonly known as soot—absorb heat in the atmosphere. For years, scientists have known that these particles are affecting Earth's warming climate, but measuring their exact effect has proved elusive.

Researchers at Michigan Technological University and Brookhaven National Laboratory, along with partners at other universities, industry, and national labs, have determined that while the shape of particles

containing black [carbon](#) does have some effect on atmospheric warming, it's important to account for the structural differences in soot particles, as well as how the particles interact with other organic and inorganic materials that coat black carbon as it travels through the atmosphere.

Published today in the *Proceedings of the National Academy of Sciences*, the article provides a framework that reconciles [model simulations](#) with laboratory and empirical observations, and that can be used to improve estimates of black carbon's impact on climate.

Every Black Carbon Particle is Unique

Black carbon's absorption of solar radiation is comparable to that of [carbon dioxide](#). Yet black carbon only remains in the atmosphere for days to weeks, while carbon dioxide can remain in the atmosphere for hundreds of years.

Scientists for years have approximated black carbon particles as spherically shaped in models that frequently became coated by other organic materials. The thought was that as the soot particles travel through the atmosphere, the coating had what is called a "lensing effect"; the coat focuses light down on the black carbon, causing increased radiation absorption. And while [soot particles](#) are indeed coated in organic materials, that coating is not uniform from particle to particle.

"When you take an image under the microscope, the particles never look perfectly like a sphere with the same coating," said Claudio Mazzoleni, professor of physics at Michigan Tech and one of the article's co-authors. "If you do a numerical calculation about perfect spheres coated by a shell, a [model](#) will show an enhanced absorption of the black carbon particles by a factor of up to three."

Empirical studies of black carbon particles demonstrate that absorption

is much less than models would suggest, calling into question the effectiveness of the model as well as our understanding of black carbon's climate forcing effect.

Research suggests that the [organic material](#) coating is not fully spherical; depending on how the organic materials cling to a black carbon particle, the resulting shape can cause the particle to act very differently even if it has the same amount of material as another soot particle that is differently shaped. But even more important is that the amount of coating might change disparately from particle to particle. These two attributes both decrease the expected absorption enhancement.

Laura Fierce, an associate atmospheric scientist at Brookhaven National Laboratory, applied the particle-resolved model to account for particle heterogeneity while modeling black carbon.

"Whereas most aerosol models simplify the representation of particle composition, the particle-resolved model tracks the composition of individual particles as they evolve in the atmosphere," Fierce said. "This model is uniquely suited to evaluate error resulting from common approximations applied in global-scale aerosol models."

Less Effect on Climate Warming Than We Thought

Essentially, the researchers have introduced into climate modeling the diversity of organic and inorganic coating on particles and the non-uniform nature of the particles themselves. By combining an empirical model with laboratory measurements, the model predicted a much lower enhancement increase in absorption by black carbon than previously thought. The updated modeling also brings the model's output much closer to what was measured in the field.

"People think black carbon has a very strong warming effect on the

atmosphere, which depends on absorption," Mazzoleni said. "If you have larger absorption, it contributes to warming and has greater climate impact. To understand how much black carbon contributes to the warming of climate, we need to understand these details because they can make a difference."

This research provides a path forward for improving predictions of black carbon's radiative effect on climate. Reducing black carbon emissions in the atmosphere can help reduce some of the effects of [climate](#) change. The results of this study suggest that a particle's [absorption](#) per mass is lower than previously thought, but how much [black carbon](#) is forcing atmospheric warming also depends on emissions levels, interactions with clouds and the distance a particle travels. And while reducing sooty emissions is significant, reducing carbon dioxide in the [atmosphere](#) is still of utmost importance.

More information: Laura Fierce et al. Radiative absorption enhancements by black carbon controlled by particle-to-particle heterogeneity in composition, *Proceedings of the National Academy of Sciences* (2020). [DOI: 10.1073/pnas.1919723117](https://doi.org/10.1073/pnas.1919723117)

Provided by Michigan Technological University

Citation: Like snowflakes, soot particles are unique, affecting climate modeling (2020, February 27) retrieved 27 April 2024 from <https://phys.org/news/2020-02-snowflakes-soot-particles-unique-affecting.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--