


Synthesizing a superatom: Opening doors to their use as substitutes for elemental atoms

February 25 2020



Based on the dendrimer template method, clusters consisting of 3, 12, 13 and other number of atoms have been synthesized. The halogen-like superatomic nature of Ga13 was structurally and electro-chemically observed as completely different to the other clusters. The results for these gallium clusters have provided candidates for superatoms. Credit: Tokyo Tech

'Superatom' is a name given to a cluster of atoms that seem to exhibit properties similar to elemental atoms. Scientists have shown particular



interest in superatomic structures, since they can be linked with atoms to produce molecules, and potentially be used to substitute certain elements in many applications.

But for superatoms to be effectively utilized, they must be specially tailored to resemble the characteristics of the corresponding elements. This transformation depends on the specific combination of electrons used. For example, when an aluminum atom with 3 valence electrons (outer shell electrons that can contribute to the formation of chemical bonds) is added to the superatom of aluminium-13, the properties change to those of a superatom of aluminium-14. Due to this modifiability of superatoms, investigating them and understanding them further is important. But previous research has been mainly theoretical, and largely focused on single clusters. Research has also not been able to synthesize superatomic clusters with sufficient volume and stability for practical application.

In a recent study, scientists from Tokyo Tech and ERATO Japan Science and Technology, led by Dr. Tetsuya Kambe and Prof Kimihisa Yamamoto, fabricated clusters of the element gallium (Ga) in solution to demonstrate the effects of changing the number of atoms in a <u>cluster</u> on the properties of the cluster. The team synthesized Ga clusters of 3, 12, 13 and other numbers of atoms using a specialized superatom synthesizer. To characterize and analyze the structural differences among the synthesized clusters, transmission electron microscopic images were captured and calculations were performed using computation tools.

The mass spectrometry revealed that the 13- and 3-atom clusters had superatomic periodicity. The 13-atom cluster differed from the other clusters structurally and electrochemically. But the 3-atom cluster with hydrogen (Ga_3H_2) was reduced to Ga_3H_2 - and was not detected, suggesting a low stability of this cluster when synthesized in the solution



medium.

The ability to alter the clusters reinforces the concept that structural change can be induced in superatoms. Describing the implications of their findings, the scientists explain: "These series of results demonstrate that it is possible to change the <u>valence electrons</u> in superatomic clusters in solution by controlling the number of constituent atoms. This in turn enables the designing and preparation of superatoms."

This study paves the way for future research to investigate the use of superatoms as substitutes for elements. As Dr. Kambe, Prof Yamamoto and team reiterate, "the <u>superatom</u> reveals an attractive strategy for creating new building blocks through the use of cluster structures."

More information: Tetsuya Kambe et al, Superatomic Gallium Clusters in Dendrimers: Unique Rigidity and Reactivity Depending on their Atomicity, *Advanced Materials* (2020). <u>DOI:</u> <u>10.1002/adma.201907167</u>

Provided by Tokyo Institute of Technology

Citation: Synthesizing a superatom: Opening doors to their use as substitutes for elemental atoms (2020, February 25) retrieved 24 April 2024 from <u>https://phys.org/news/2020-02-superatom-doors-substitutes-elemental-atoms.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.