


Team identifies low-energy solar particles from beyond Earth near the Sun

February 3 2020



Using data from NASA's Parker Solar Probe, an SwRI-led team identified low-energy particles, the smoking gun pointing to interactions between slow- and fast-moving regions of the solar wind accelerating high-energy particles from beyond the orbit of Earth. Using Integrated Science Investigation of the Sun (ISIS) instrument data, they measured low-energy particles in the near-Sun environment that had likely traveled back toward the Sun, slowing against the tide of the solar wind while still retaining surprising energies. Credit: NASA/Johns Hopkins APL/Steve Gribben

Using data from NASA's Parker Solar Probe (PSP), a team led by Southwest Research Institute identified low-energy particles lurking near the Sun that likely originated from solar wind interactions well beyond Earth orbit. PSP is venturing closer to the Sun than any previous probe, carrying hardware SwRI helped develop. Scientists are probing the enigmatic features of the Sun to answer many questions, including how to protect space travelers and technology from the radiation associated with solar events.

"Our main goal is to determine the acceleration mechanisms that create and transport dangerous high-energy particles from the solar atmosphere into the solar system, including the near-Earth environment," said Dr. Mihir Desai, a mission co-investigator on the Integrated Science Investigation of the Sun (IS \odot IS) instrument suite, a multi-institutional project led by Principal Investigator Prof. Dave McComas of Princeton University.. IS \odot IS consists of two instruments, Energetic Particle Instrument-High (EPI-Hi) and Energetic Particle Instrument-Low (EPI-Lo). "With EPI-Lo, we were able to measure extremely low-energy particles unexpectedly close to the solar environment. We considered many explanations for their presence, but ultimately determined they are the smoking gun pointing to interactions between slow- and fast-moving regions of the solar wind that accelerate high-energy particles from beyond the orbit of Earth. Some of those travel back toward the Sun, slowing against the tide of the outpouring solar wind but still retaining surprisingly high energies."

PSP, which will travel within 4 million miles of the Sun's surface, is collecting new solar data to help scientists understand how [solar events](#), such as coronal mass ejections, impact life on Earth. During the rising portion of the Sun's activity cycle, our star releases huge quantities of energized matter, magnetic fields and electromagnetic radiation in the form of coronal mass ejections (CMEs). This material is integrated into the solar wind, the steady stream of charged particles released from the

Sun's upper atmosphere. The high-energy solar energetic particles (SEPs) present a serious radiation threat to human explorers living and working outside low-Earth orbit and to technological assets such as communications and scientific satellites in space. The mission is making the first-ever direct measurements of both the low-energy source populations as well as the more hazardous, higher-energy particles in the near-Sun environment, where the acceleration takes place.

When the Sun's activity reaches a lull, roughly about every 11 years, solar equatorial regions emit slower solar wind streams, traveling around 1 million miles per hour, while the poles spew faster streams, traveling twice as fast at 2 million miles per hour. Stream Interaction Regions (SIRs) are created by interactions at boundaries between the fast and slow solar wind. Fast-moving streams tend to overtake slower streams that originate westward of them on the Sun, forming turbulent corotating interaction regions (CIRs) that produce shock waves and accelerated particles, not unlike those produced by CMEs.

"For the first time, we observed low-energy particles from these CIRs near the orbit of Mercury," Desai said. "We also compared the PSP data with data from STEREO, another solar energy probe. By measuring the full range of energetic populations and correlating the data with other measurements, we hope to get a clear picture of the origin and the processes that accelerate these particles. Our next step is to integrate the data into models to better understand the origin of SEPs and other materials. Parker Solar Probe will solve many puzzling scientific questions—and is guaranteed to generate new ones as well."

This research is described in the paper "Properties of Suprathermal-through-Energetic He Ions Associated with Stream interaction regions Observed over Parker Solar Probe's First Two Orbits," published February 3 in a special issue of the *Astrophysical Journal Supplement Series* devoted exclusively to the first science results from the Parker

Solar Probe mission.

More information: M. I. Desai et al, Properties of Suprathermal-through-energetic He Ions Associated with Stream Interaction Regions Observed over the Parker Solar Probe's First Two Orbits, *The Astrophysical Journal Supplement Series* (2020). [DOI: 10.3847/1538-4365/ab65ef](https://doi.org/10.3847/1538-4365/ab65ef)

Provided by Southwest Research Institute

Citation: Team identifies low-energy solar particles from beyond Earth near the Sun (2020, February 3) retrieved 10 April 2024 from <https://phys.org/news/2020-02-team-low-energy-solar-particles-earth.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--