


Researchers realize two-photon pumped nanolaser from formamidinium perovskites

February 14 2020, by Zhang Nannan



Tunable photoluminescence and lasing spectra from FAPbX₃ NCs. Credit: SIOM

Formamidinium (FA) perovskites have exhibited outstanding optoelectronic properties in efficient solar cells and light-emitting diodes. However, their development on nanolaser application has rarely been explored, especially the up-conversion lasing performance.

Recently, a collaborative research team from Shanghai Institute of

Optics and Fine Mechanics (SIOM) of the Chinese Academy of Sciences (CAS) and Chongqing University has succeeded in obtaining two-photon pumped lasing from colloidal FA-perovskite nanocrystals (NCs) at [room temperature](#). This work was published in *ACS Photonics*.

In the experiment, researcher prepared the FA-perovskite NCs as gain medium by a simple ligand-assisted reprecipitation strategy. These colloidal FAPbX₃ NCs exhibited tunable bright emission (from 433 to 667 nm) through halide exchange.

Simultaneously, by tuning the ratio of the surface organic ligands, the morphology of FAPbBr₃ NCs could be transferred from [quantum dots](#) to nanoplatelets, and consequently, the emission spectra were also tuned in 520-542 nm.

Subsequently, they demonstrated the excellent nonlinear characteristic of NCs. A large two-photon absorption coefficient (0.76 cm/GW) and high optical net gain (480 cm⁻¹) were achieved, suggesting that FAPbX₃ NCs could be regarded as an excellent optical gain medium.

Furthermore, the low-threshold amplified [spontaneous emission](#) (ASE) under both two- and one-photon excitations was observed. The temperature-dependent ASE action revealed the strong temperature-tolerant ASE action with a high characteristic temperature of 308 K, indicating the weak requirements of heat management of FA-based devices.

Finally, FAPbBr₃ NCs were coupled into a hollow capillary tube. Stable two-photon excited whispering-gallery-mode lasing was successfully achieved with a low threshold of $\approx 310 \mu\text{J}/\text{cm}^2$.

These findings suggest that FAPbX₃ NCs can act as promising gain media for high-performance upconverted nanolasers toward

optoelectronic application.

More information: Zhengzheng Liu et al. Two-Photon Pumped Amplified Spontaneous Emission and Lasing from Formamidinium Lead Bromine Nanocrystals, *ACS Photonics* (2019). [DOI: 10.1021/acsphotonics.9b01226](https://doi.org/10.1021/acsphotonics.9b01226)

Provided by Chinese Academy of Sciences

Citation: Researchers realize two-photon pumped nanolaser from formamidinium perovskites (2020, February 14) retrieved 18 April 2024 from <https://phys.org/news/2020-02-two-photon-nanolaser-formamidinium-perovskites.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.