


# Air pollution declines in Spain after implementation of measures to fight COVID-19

March 25 2020

Satellite data confirm a decrease in the nitrogen dioxide emissions during the COVID-19 pandemic in Spain



Credit: Universitat Politècnica de València

The concentration levels of nitrogen dioxide, a primary air contaminant, have declined 64% in major Spanish cities after the implementation of measures to fight COVID-19. Levels have fallen most in Barcelona, with an 83% decline. In Madrid, levels have dropped 73% and 64% in

València.

These are some of the results of a study carried out by researchers from the Universitat Politècnica de València (UPV), belonging to the Centro de Tecnologías Físicas (Physical Technology Center). To carry out this study, the UPV team analyzed satellite images of the Sentinel-5P mission of the Copernicus program of the European Space Agency (ESA), comparing data from March 10-14 against data from March 15-20. Based on this data, they generated a series of maps that show the concentrations of [nitrogen](#) dioxide in major Spanish cities.

According to Elena Sánchez-García, researcher at the Land and Atmosphere Remote Sensing (LARS-UPV) group of the UPV's Centro de Tecnologías Físicas, nitrogen dioxide is an important indicator of air quality. "High concentrations of nitrogen can affect the respiratory system and aggravate certain medical conditions. In addition, this gas is related to the formation of acid rain," she said.

Nitrogen oxides like NO<sub>2</sub> in the urban atmosphere originate from combustion reactions at high temperatures that are mainly produced from motorized vehicles. Oxygen and nitrogen are combined to make [nitric oxide](#) (NO), which is later partially oxidized, creating nitrogen dioxide.

"As has been verified in the case of Wuhan (China), and northern Italy, our study shows how the confinement measures and reduction of economic activity have translated into a clear decrease in air pollution throughout the country," said Elena Sánchez-García.

## **Additional Data**

In addition to València, Madrid and Barcelona, the study also took data from seven other cities. In the Autonomous Community of Valencia,

[nitrogen dioxide](#) levels have dropped most in Castellón, with a 76% decline. Of the cities analyzed, it had the second-largest reduction, behind only Barcelona. In Alicante, the decline was 68%. Though [atmospheric variability](#) (winds and precipitation) can affect the numbers calculated for each [city](#), the effect of the current situation of confinement is predominant. Of the other cities studied, in Bilbao, the contamination from one period to the other dropped 66%; in Gijón, 65%; in Málaga, 55%; in Zaragoza, 52%; and in Sevilla, 36%.

The study is framed within the research carried out by the LARS-UPV group of the UPV, which featured the participation of Elena Sánchez García, Itziar Irakulis Loitxate and Luis Guanter. Their work is focused on the development of remote detection techniques for globally monitoring the emissions of gases into the atmosphere.

Provided by Universitat Politècnica de València

Citation: Air pollution declines in Spain after implementation of measures to fight COVID-19 (2020, March 25) retrieved 15 May 2024 from <https://phys.org/news/2020-03-air-pollution-declines-spain-covid-.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--