

# How plants sound the alarm about danger

March 13 2020

---



*Arabidopsis thaliana*, a small flowering plant in the mustard family. Credit: Salk Institute

Just like humans and other animals, plants have hormones. One role of plant hormones is to perceive trouble—whether an insect attack, drought or intense heat or cold—and then signal to the rest of the plant to respond.

A multicenter team led by current and former investigators from the Salk Institute is reporting new details about how plants respond to a hormone called [jasmonic acid](#), or jasmonate. The findings, which were published in *Nature Plants* on March 13, 2020, reveal a complex communication network. This knowledge could help researchers, such as members of Salk's Harnessing Plants Initiative, develop crops that are hardier and more able to withstand assault, especially in an era of rapid climate change.

"This research gives us a really detailed picture of how this hormone, jasmonic acid, acts at many different levels," says Professor Joseph Ecker, co-corresponding author and Howard Hughes Medical Institute investigator. "It enables us to understand how environmental information and developmental information is processed, and how it ensures proper growth and development."

The plant used in the study was *Arabidopsis thaliana*, a small flowering plant in the mustard family. Because its genome has been well characterized, this plant is a popular model system. Scientists can take what they learn in *A. thaliana* and apply it to other plants, including those grown for food. Jasmonic acid is found not only in *A. thaliana* but throughout the plant kingdom.

"Jasmonic acid is particularly important for a plant's defense response against fungi and insects," says co-first author Mark Zander, a staff researcher in Ecker's lab. "We wanted to precisely understand what

happens after jasmonic acid is perceived by the plant. Which genes are activated and deactivated, which proteins are produced and which factors are in control of these well-orchestrated cellular processes?"

The researchers started with [plant seeds](#) grown in petri dishes. They kept the seeds in the dark for three days to mimic the first few days of a seed's life, when it is still underground. "We know this growth stage is super important," says co-first author and co-corresponding author Mathew Lewsey, an associate professor at La Trobe University in Melbourne, Australia, who previously worked in Ecker's lab. The first few days in the soil are a challenging time for seedlings, as they face attacks from insects and fungi. "If your seeds don't germinate and successfully emerge from the soil, then you will have no crop," Lewsey adds.

After three days, the [plants](#) were exposed to jasmonic acid. The researchers then extracted the DNA and proteins from the plant cells and employed specific antibodies against their proteins of interest to capture the exact genomic location of these regulators. By using various computational approaches, the team was then able to identify genes that are important for the plant's response to jasmonic acid and, moreover, for the cellular cross-communication with other plant hormone pathways.

Two genes that rose to the top in their degree of importance across the system were MYC2 and MYC3. These genes code for proteins that are [transcription factors](#), which means that they regulate the activity of many other genes—or thousands of other genes in this case.

"In the past, the MYC genes and other transcription factors have been studied in a very linear fashion," Lewsey explains. "Scientists look at how one gene is connected to the next gene, and the next one, and so on. This method is inherently slow because there are a lot of genes and lots

of connections. What we've done here is to create a framework by which we can analyze many genes at once."

"By deciphering all of these gene networks and subnetworks, it helps us to understand the architecture of the whole system," Zander says. "We now have this very comprehensive picture of which genes are turned on and off during a plant's defense response. With the availability of CRISPR gene editing, these kinds of details can be useful for breeding crops that are able to better withstand attacks from pests."

Another noteworthy aspect of this work is that all of the data from the research has been made available on Salk's website. Researchers can use the site to search for more information about [genes](#) they study and find ways to target them.

**More information:** Zander, M., et al. Integrated multi-omics framework of the plant response to jasmonic acid. *Nat. Plants* 6, 290–302 (2020). [doi.org/10.1038/s41477-020-0605-7](https://doi.org/10.1038/s41477-020-0605-7)

Provided by Salk Institute

Citation: How plants sound the alarm about danger (2020, March 13) retrieved 23 June 2024 from <https://phys.org/news/2020-03-alarm-danger.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.