

Amber specimens reveal origin of long mouthpart of scorpionflies

March 4 2020



Ecological reconstruction of Mesozoic Aneuretopsyhidae. Credit: NIGPAS

An international research group led by Prof. Wang Bo from the Nanjing Institute of Geology and Palaeontology of the Chinese Academy of Sciences (NIGPAS) has found a new genus, including two new aneuretopsyhid species from early Late Cretaceous (99 million years ago) Burmese amber, which reveals new anatomically significant details of the elongate mouthpart elements.


Mesopsychoid scorpionflies are peculiar Mesozoic insects with a distinctly elongate mouthpart and are considered to be a critical group of pollinators prior to the rise of angiosperms.

A new genus found from 99-million-year-old Burmese amber reveals the origin of scorpionflies' long mouthpart. This discovery was reported in *Science Advances* on March 4. Aneuretopsyhidae is a family of mecopteran insects with a long siphonate mouthpart. In particular, this family is the key to understanding both the [early evolution](#) of highly modified mouthparts in Mesopsychoida and arguably the origin of fleas.

Previously, all known aneuretopsyhids were from compression fossils, and the detailed structure of their mouthparts was still unclear.

Now, however, an international research group led by Prof. Wang Bo from the Nanjing Institute of Geology and Palaeontology of the Chinese Academy of Sciences (NIGPAS) has found a [new genus](#), including two new aneuretopsyhid species from early Late Cretaceous (99 million years ago) Burmese amber, which reveals new anatomically significant details of the elongate mouthpart elements.

The aneuretopsychid mouthpart in the new amber fossils consists of one pair of galeae and one unpaired central hypopharynx. During feeding, the galeae would come together temporarily and enclose the hypopharynx thus forming a functional tube.




Aneuretopsychidae from Late Cretaceous Burmese amber. Credit: NIGPAS

The structures of the new three-dimensionally preserved fossils thus reveal that the aneuretopsychid mouthpart is not labial but maxillary in origin.

The phylogenetic results based on 38 taxa and 54 discrete characters support the monophyly of Mesopsuchoidea and demonstrate that an elongate mouthpart is one of its key synapomorphies, challenging the view that the long-proboscid condition independently originated two or three times in this clade.

In addition, the mouthpart of Mesopsuchoidea differs structurally from the highly modified piercing mouthparts of Siphonaptera. So, neither Aneuretopsychidae nor Mesopsuchoidea is a sister group to Siphonaptera.



Wing venation of Aneuretopsychidae. Credit: NIGPAS

In the Burmese amber forest, at least five families of long-proboscid insects have been discovered, further revealing the variety and complexity of mid-Cretaceous pollinating insects.

This study provides new insights into the separate origin of the long mouthpart of Mesopsychosidea and fleas, and the evolution of Cretaceous pollinating insects.

More information: X. Zhao et al., "Mouthpart homologies and life habits of Mesozoic long-proboscid scorpionflies," *Science Advances* (2020). DOI: [10.1126/sciadv.aay1259](https://doi.org/10.1126/sciadv.aay1259) , advances.sciencemag.org/content/6/10/eaay1259

Provided by Chinese Academy of Sciences

Citation: Amber specimens reveal origin of long mouthpart of scorpionflies (2020, March 4)
retrieved 3 May 2024 from

<https://phys.org/news/2020-03-amber-specimens-reveal-mouthpart-scorpionflies.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--