

# Defaunation in rainforests could have more severe consequences than reported

March 31 2020

---



The crimson-rumped toucanet (*Aulacorhynchus haematopygus*) is one of the large-bodied birds inhabiting the rainforest. Credit: Maximilian Vollstädt

Tropical rainforests are emptying out due to the ongoing extinction of animal species caused by overhunting and forest fragmentation. But not only species as such, also the number of individuals of species living in

rainforests continue to decline. Large-bodied species, which are particularly vulnerable to changes, decline at a comparably faster rate than other species. A new study by researchers at the Senckenberg Biodiversity and Climate Research Centre shows that the resulting animal downsizing in rainforests might have severe consequences for ecosystem services.

"The effect of downsizing of animal communities is three times larger for services that are provided by ecological networks than for their structural stability. This is because the [species](#) that go extinct first are those who contribute unique services and keep the [rainforest](#) function," says researcher Dr. Isabel Donoso, an Alexander-von-Humboldt fellow who led the study at Senckenberg.

Donoso and her team simulated the effect of the loss of large-bodied fruit-eating birds for eight locations in the Andes. The simulation was based on data about interactions between bird and [plant species](#) recorded over many years. To measure the [ecosystem services](#) provided by birds, the researchers simulated the distance of avian seed dispersal. Seed dispersal by animals is a pivotal service in rainforests as most tropical trees require animals to disperse their seeds. The study is one of first attempts to quantify the consequences of animal downsizing for the functioning of rainforest ecosystems. Press images may be used at no cost for editorial reporting, provided that the original author's name is published, as well. The images may only be passed on to third parties in the context of current reporting.

"If the largest 10 percent of the fruit-eating birds were to go locally extinct, the distance over which seeds would be dispersed would shrink by almost 40 percent. In contrast to this, the structural stability of an ecological network would be changed to a much lesser degree. A 10 percent loss of the largest fruit-eating birds would reduce the number of interactions between fruit-eating birds and plants by only 10 percent,"

says Dr. Matthias Schleuning, Senckenberg Biodiversity and Climate Research Centre explains the results of the simulation.

So why should one care if in the future seeds of rainforest trees are dispersed short-distance rather than long-distance? First of all, long-distance [seed dispersal](#) helps the genetic exchange between forest patches and makes them more resilient to change. What's more important: the nearer a [seed](#) remains to the parent tree, it becomes less likely that it will germinate and grow. Because large-bodied fruit-eating birds predominately disperse large seeds, tree with large seeds are the ones supposedly losing out.

"Future rainforests will probably look very different in terms of tree composition than the current ones. Other studies have already shown that such future rainforests will likely be able to store much less carbon," explains Donoso. In light of this it would be short-sighted to underestimate the looming consequences of animal downsizing, warns Schleuning: "We need to ramp up our efforts to protect large-bodied [animals](#) in rainforests. Only then global climate and we all will continue to benefit from the services [tropical rainforests](#) provide to us."

**More information:** Isabel Donoso et al. Downsizing of animal communities triggers stronger functional than structural decay in seed-dispersal networks, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-15438-y](https://doi.org/10.1038/s41467-020-15438-y)

Provided by Senckenberg Research Institute and Natural History Museum

Citation: Defaunation in rainforests could have more severe consequences than reported (2020, March 31) retrieved 20 May 2024 from <https://phys.org/news/2020-03-defaunation-rainforests->

[severe-consequences.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.