

New feathered dinosaur was one of the last surviving raptors

March 26 2020



A new feathered dinosaur that lived in New Mexico 67 million years ago is one of the last known surviving raptor species, according to a new publication in the journal *Scientific Reports*. *Dineobellator notohesperus* adds to scientists' understanding of the paleo-biodiversity of the American Southwest, offering a clearer picture of what life was like in this region near the end of the reign of the dinosaurs. Credit: Sergey Krasovskiy

A new feathered dinosaur that lived in New Mexico 67 million years ago is one of the last known surviving raptor species, according to a new publication in the journal *Scientific Reports*.

Dineobellator notohesperus adds to scientists' understanding of the paleobiodiversity of the American Southwest, offering a clearer picture of what life was like in this region near the end of the reign of the dinosaurs.

Steven Jasinski, who recently completed his Ph.D. in Penn's Department of Earth and Environmental Sciences in the School of Arts and Sciences, led the work to describe the new species, collaborating with doctoral advisor Peter Dodson of the School of Veterinary Medicine and Penn Arts and Sciences and as well as Robert Sullivan of the New Mexico Museum of Natural History and Science in Albuquerque.

In 2008, Sullivan found fossils of the new species in Cretaceous rocks of the San Juan Basin, New Mexico. He, along with his field team of Jasinski and James Nikas, collected the specimen on U.S. federal land under a permit issued by the Bureau of Land Management. The entire specimen was recovered over four field seasons. Jasinski and his coauthors gave the species its official name, *Dineobellator notohesperus*, which means "Navajo warrior from the Southwest," in honor of the people who today live in the same region where this dinosaur once dwelled.

Dineobellator, as well as its Asian cousin *Velociraptor*, belong to a group of dinosaurs known as the dromaeosaurids. Members of this group are commonly referred to as "raptor" dinosaurs, thanks to movies such as "Jurassic Park" and "Jurassic World." But unlike the terrifying beasts depicted in film, *Dineobellator* stood only about 3.5 feet (about 1 meter) at the hip and was 6 to 7 feet (about 2 meters) long—much smaller than its Hollywood counterparts.

Raptor dinosaurs are generally small, lightly built predators. Consequently, their remains are rare, particularly from the southwestern United States and Mexico. "While dromaeosaurids are better known

from places like the northern United States, Canada, and Asia, little is known of the group farther south in North America," says Jasinski.

While not all of the bones of this dinosaur were recovered, bones from the forearm have quill nobs—small bumps on the surface where feathers would be anchored by ligaments—an indication that *Dineobellator* bore feathers in life, similar to those inferred for *Velociraptor*.

Features of the animal's forelimbs, including enlarged areas of the claws, suggest this dinosaur could strongly flex its arms and hands. This ability may have been useful for holding on to prey—using its hands for smaller animals such as birds and lizards, or perhaps its arms and feet for larger species such as other dinosaurs.

Its tail also possessed unique characteristics. While most raptors' tails were straight and stiffened with rod-like structures, *Dineobellator*'s tail was rather flexible at its base, allowing the rest of the tail to remain stiff and act like a rudder.

"Think of what happens with a cat's tail as it is running," says Jasinski.

"While the tail itself remains straight, it is also whipping around constantly as the animal is changing direction. A stiff tail that is highly mobile at its base allows for increased agility and changes in direction, and potentially aided *Dineobellator* in pursuing prey, especially in more open habitats."

This new dinosaur provides a clearer picture of the biology of North American dromaeosaurid dinosaurs, especially concerning the distribution of feathers among its members.

"As we find evidence of more members possessing feathers, we believe it is likely that all the dromaeosaurids had feathers," says Jasinski. The discovery also hints at some of the predatory habits of a group of iconic

meat-eating dinosaurs that lived just before the extinction event that killed off all the dinosaurs that weren't birds.

Jasinski plans to continue his field research in New Mexico with the hope of finding more fossils.

"It was with a lot of searching and a bit of luck that this dinosaur was found weathering out of a small hillside," he says. "We do so much hiking and it is easy to overlook something or simply walk on the wrong side of a hill and miss something. We hope that the more we search, the better chance we have of finding more of *Dineobellator* or the other [dinosaurs](#) it lived alongside."

More information: New Dromaeosaurid Dinosaur (Theropoda, Dromaeosauridae) from New Mexico and Biodiversity of Dromaeosaurids at the end of the Cretaceous, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-61480-7](https://doi.org/10.1038/s41598-020-61480-7) , www.nature.com/articles/s41598-020-61480-7

Provided by University of Pennsylvania

Citation: New feathered dinosaur was one of the last surviving raptors (2020, March 26) retrieved 27 April 2024 from <https://phys.org/news/2020-03-feathered-dinosaur-surviving-raptors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
