

Feeding fusion: hydrogen ice pellets prove effective for fueling fusion plasmas

March 10 2020, by John Greenwald



PPPL physicist Oak Nelson. Credit: Elle Starkman/PPPL Office of Communications.

Researchers have found that injecting pellets of hydrogen ice rather than puffing hydrogen gas improves fusion performance at the DIII-D



National Fusion Facility, which General Atomics operates for the U.S. Department of Energy (DOE). The studies by physicists based at DOE's Princeton Plasma Physics Laboratory (PPPL) and Oak Ridge National Laboratory (ORNL) compared the two methods, looking ahead to the fueling that will be used in ITER, the international fusion experiment under construction in France.

Improve the temperature

The researchers showed that icy pellets of <u>hydrogen</u> improve the temperature of the <u>fusion</u> plasma when compared with the gas fueling method now typically used in doughnut-shaped fusion facilities called tokamaks. Higher temperatures are beneficial for the fusion reactions. The results on DIII-D are encouraging for ITER, which plans to use pellet injection to fuel its hot inner core.

Fusion, the power that drives the sun and stars, combines light elements in the form of plasma—the state of matter that consists of positively charged atomic nuclei and negatively charged electrons—to create massive amounts of energy. Scientists seek to replicate fusion on Earth for a safe, clean and virtually inexhaustible supply of power to generate electricity.

One challenge for producing fusion energy is how to get cold hydrogen fuel into the hot plasma core. The sun has all the hydrogen that it needs for billions of years, but fusion reactors on Earth must constantly feed hydrogen into the plasma to sustain the fusion reactions. Puffing roomtemperature gas is the most common way to inject hydrogen in current experiments.

Bigger and hotter



However, as fusion reactors get bigger and hotter it will become harder for the gas to penetrate into the core of the reactor where fusion reactions take place. New methods thus need to be developed to feed the fusion core without degrading the plasma performance.

The joint research effort on DIII-D compared the two fueling methods in high-performance plasmas planned for ITER. The experiments revealed a significantly higher pressure of plasma—a key to fusion reactions—using hydrogen ice compared to gas injection when the rate of fueling is roughly evenly matched between the two methods.

"The fueling plays a big role in the edge <u>plasma</u> performance," said Andrew "Oak" Nelson, a graduate student in the Program in Plasma Physics at Princeton University and first author of the Nuclear Fusion <u>article describing these results</u>. Nelson is part of a multi-institutional team that carefully designed and executed the experiments.

Scientists at ORNL

The technology for injecting the ice pellets was developed by scientists at ORNL. Interpretation of the experimental results requires sophisticated scientific instruments developed by multiple collaborating institutions on DIII-D. "It's great to see how our multi-institutional effort came together to tackle this important fueling question for ITER and future reactors," said Morgan Shafer, a lead research scientist at ORNL and a coauthor of the paper.

The research also demonstrates how graduate students can make important contributions to fusion energy by working on these large national research facilities. "For a <u>graduate student</u> to play an important role in this experimental study on DIII-D is impressive," said Egemen Kolemen, a PPPL and Princeton University physicist who was an advisor for the project. "Oak's success shows how large fusion experiments



provide significant leadership opportunities for students and early career scientists."

More information: A.O. Nelson et al, Setting the H-mode pedestal structure: variations of particle source location using gas puff and pellet fueling, *Nuclear Fusion* (2019). DOI: 10.1088/1741-4326/ab5e65

Provided by Princeton Plasma Physics Laboratory

Citation: Feeding fusion: hydrogen ice pellets prove effective for fueling fusion plasmas (2020, March 10) retrieved 29 April 2024 from <u>https://phys.org/news/2020-03-fusion-hydrogen-ice-pellets-effective.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.