


When rocks lay the groundwork for the origin of life

March 2 2020, by University Of Strasbourg



The diagram illustrates the chemical pathways for the synthesis of key biomolecules from hydrothermal minerals and gases. The pathways are as follows:

- Pyruvate Pathway:** $\text{CO}_2 + \text{H}_2 \rightarrow \text{Pyruvate}$
- Acetate Pathway:** $\text{CO}_2 + \text{H}_2 \rightarrow \text{Acetate}$
- Methane Pathway:** $\text{CO}_2 + \text{H}_2 \rightarrow \text{Methane (CH}_4\text{)}$
- Formate Pathway:** $\text{CO}_2 + \text{H}_2 \rightarrow \text{Formate} \rightarrow \text{H}_3\text{C} \rightarrow \text{CO} \rightarrow \text{Acetate} \rightarrow \text{Pyruvate}$
- Alternative Pathway:** $\text{CO}_2 + \text{H}_2 \rightarrow \text{CO} \rightarrow \text{H}_3\text{C} \rightarrow \text{Methane}$

Hydrothermal minerals are shown as the source of CO_2 and H_2 for these reactions.

Susan Lang, U. of SC. / NSF / ROV Jason / 2018
© Woods Hole Oceanographic Institution

Mineral catalysts found in deep-sea vents convert carbon dioxide and hydrogen to biomolecules, showing striking parallels to known biological pathways. Credit: Susan Lang, U. of SC. / NSF / ROV Jason / 2018 © Woods Hole Oceanographic Institution

Mineral catalysts found in deep-sea vents convert CO₂ and H₂ to biomolecules, showing striking parallels to known biological pathways.

An international collaboration of researchers in Germany, France and Japan investigated the catalytic activities of minerals found in deep-sea hydrothermal vents. The results suggest that mineral-driven [chemical reactions](#) might be closely mapped onto microbial carbon metabolism.

Life on Earth could not have started without metabolism. Alkaline hydrothermal vents have been proposed as a possible environment where a primitive form of metabolism (protometabolism) predating cellular life could have emerged. The first steps of metabolism, which would have started as geological chemistry, might still be conserved in biochemical pathways that [evolutionary biologists](#) have identified as being ancient. One of them is the *acetyl-CoA pathway*. Microorganisms use it to convert carbon dioxide (CO₂) to acetate and pyruvate, serving as an entry point to a more complex carbon metabolism. Interestingly, it is the only known biological carbon fixation route that uses hydrogen (H₂) as a source of electrons, and H₂ is very abundant in alkaline hydrothermal vent systems. Still, geochemically realistic CO₂ fixation leading to higher-carbon biomolecules under biologically relevant deep-sea vent conditions has been challenging.

Looking for a bridge between the abiotic (non-life) and biotic (life), the researchers in Germany, France and Japan became inspired by the characteristics of the biological acetyl-CoA pathway. In biology,

complex enzymes facilitate the reaction between CO_2 and H_2 , but hydrothermal minerals can serve as catalysts for the non-enzymatic version of the same reaction. Prof. Joseph Moran, an organic chemist interested in the protometabolic origin of life and leader of the Laboratory of Chemical Catalysis at the University of Strasbourg (France) explains: "What acetyl-CoA pathway enzymes and hydrothermal minerals have in common is transition metals like iron (Fe) and nickel (Ni). These metals are also commonly used as catalysts in organic chemistry to promote notoriously difficult reactions." The Strasbourg-based team had already explored fixing CO_2 alone using metallic iron in the origin of life context. It was the right time to move on to geologically abundant mineral catalysts.

Biologist Prof. William Martin from the Heinrich Heine University in Düsseldorf has been investigating the parallels between biochemical and geochemical CO_2 fixation for two decades. "Under physiological conditions, H_2 alone is not reducing enough to fix CO_2 . We wanted to find catalysts that could have supported life at the very beginning, before the complex biological mechanisms were in place," says Martin. Together with chemist Dr. Martina Preiner (co-first author of the paper), they reached out to a research group at the Max Planck Institute for Coal Research in Mülheim, Germany, led by Dr. Harun Tüysüz, who specializes in heterogeneous catalysis. After some brainstorming, Martin, Moran and Tüysüz decided the experiments would begin with H_2 , CO_2 and minerals found in alkaline hydrothermal vents: awaruite and magnetite. However, the three teams were not the only ones with a similar goal. Researchers in Tsukuba and Sapporo (Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology) had started reducing CO_2 with H_2 over iron sulfide catalysts (e.g. greigite mineral), which are also common in deep-sea vent systems. Microbiologist Dr. Kensuke Igarashi (co-first author of the study), who performed experiments with greigite in Sapporo, explains: "It was clear we can achieve a deeper understanding of our findings if we share our

results with each other."

Indeed, this open-minded approach proved to be very fruitful. Although CO₂ fixation with H₂ over three hydrothermal minerals—awaruite (Ni₃Fe), magnetite (Fe₃O₄) and greigite (Fe₃S₄)—was performed in three different laboratories all over the world, the results lead to the same conclusion. "All these minerals catalyze the fixation of CO₂ with H₂ in water and yield products which correspond to those of the biological acetyl-CoA pathway: formate, acetate and pyruvate," says Dr. Kamila Muchowska, a Strasbourg-based chemist and co-first author of the paper, responsible for magnetite experiments. "We studied the surface of all the minerals tested and we have strong evidence the reactions are catalytic and occur on the [mineral](#) surface" explains Tüysüz. "In the hydrothermal process of CO₂ fixation hydrogen acts as an electron- and energy donor, just like in biology. It's truly astonishing," Preiner adds.

The researchers' findings suggest that simple abiotic geochemical reactions could have paved the way for early metabolism that microbes still use today. Hydrothermal minerals could have functioned as geochemical primers for the later emergence of the enzymatic acetyl-CoA pathway. These results might not only be relevant for terrestrial hydrothermal systems, and the scientists do not exclude the possibility that protometabolism could be initiated by a similar route elsewhere.

The study is published in the journal *Nature Ecology & Evolution*.

More information: A hydrogen-dependent geochemical analogue of primordial carbon and energy metabolism, *Nature Ecology & Evolution* (2020). [DOI: 10.1038/s41559-020-1125-6](https://doi.org/10.1038/s41559-020-1125-6)

Provided by University of Strasbourg

Citation: When rocks lay the groundwork for the origin of life (2020, March 2) retrieved 11 May 2024 from <https://phys.org/news/2020-03-groundwork-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.