


# New metabolism discovered in bacteria

March 30 2020



While acetic acid creating bacteria and methanogens are dependent on the transfer of hydrogen in anoxic biotopes, *Acetobacterium woodii* recycles hydrogen within its cell. Credit: Sarah Ciurus, Goethe University Frankfurt.

They make sauerkraut sour, turn milk into yogurt and cheese, and give rye bread its intensive flavour: bacteria that ferment nutrients instead of using oxygen to extract their energy. *Acetobacterium woodii* (short: A. woodii) is one of these anaerobic living microbes. Cheese and bread are not its line of business—it lives far from oxygen in the sediments on the floor of the ocean, and can also be found in sewage treatment plants and the intestines of termites.

These biotopes are teeming with microbes that use the [organic substances](#) to their advantage in different ways. A number of bacteria

ferment sugars, [fatty acids](#) and alcohols to acetic [acid](#), also creating [hydrogen](#) (H<sub>2</sub>) in the process. In higher concentration, however, hydrogen inhibits the fermentation—too much hydrogen stops the fermentation reaction. For this reason, fermenting bacteria live together with microbes that depend on precisely this hydrogen—methanogens, for example, that create methane from hydrogen and carbon dioxide and thus gain energy. Both partners profit from this association—and are simultaneously so dependent on each other that neither one can survive without the other.

*A. woodii* masters both disciplines of the anaerobic "hydrogen association": it can ferment organic substances into acetic acid, and can also form acetic acid from carbon dioxide and hydrogen. In doing so, *A. woodii* recycles the important hydrogen within its own cell, as has now been discovered by the microbiologists in Professor Volker Müller's team at the Institute for Molecular Biosciences at Goethe University Frankfurt.

In the laboratory, the Frankfurt scientists turned off the gene for the enzyme that creates hydrogen in *A. woodii*, which is called hydrogenase. The result: the bacterium was only able to grow, for example in a medium with fructose, if hydrogen was added externally. Different additional tests confirmed that both paths for creating [acetic acid](#) are connected to hydrogen that does not leave the cell.

"Though the 'hydrogen recycling' we discovered, *A. woodii* possesses a maximum of metabolic flexibility," says the Frankfurt experimenter Dr. Anja Wiechmann. "In one cycle, it can both create and use hydrogen itself, or utilise hydrogen from external sources. This makes it capable of living both from organic as well as solely from inorganic substances."

Professor Volker Müller explains: "Our findings have implications far beyond the study of *Acetobacterium woodii*. There have already been

speculations that many ancient life forms possess the kind of metabolism that we have described in *A. woodii*. This is assumed, for example for the Asgard archaea that were just discovered a few years ago on the seabed off of California. Our investigations provide the first evidence that these paths of metabolism actually exist."

**More information:** Anja Wiechmann et al, It does not always take two to tango: "Syntrophy" via hydrogen cycling in one bacterial cell, *The ISME Journal* (2020). [DOI: 10.1038/s41396-020-0627-1](https://doi.org/10.1038/s41396-020-0627-1)

Provided by Goethe University Frankfurt am Main

Citation: New metabolism discovered in bacteria (2020, March 30) retrieved 17 April 2024 from <https://phys.org/news/2020-03-metabolism-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.