

Destroying Nigeria's riverside forests is bad for the freshwater ecosystem

March 3 2020, by Emmanuel O. Akindele

Nigeria's [population growth](#) – at a rate of more than [2.5%](#) a year—is indirectly driving all kinds of environmental degradation. There is an incessant demand for more space for people to live in and use. One of the [consequences](#) is the destruction of forests. Forests that are close to water, known as riparian forests, tend to be [most vulnerable](#).

Access to a watercourse has often [informed the choice](#) of human settlement or activity. Needing a year-long supply of [water](#) for their crops, farmers often convert riparian forests to agricultural lands. In [many cases](#), especially in the dry season, such farmlands encroach almost into the waterways.

Other kinds of business also need a source of water. In Nigeria, car-wash centres are often sited along rivers and streams where forests once existed. Block-making operations, too, are often situated close to waterways.

The rapid decline in freshwater biodiversity has been [attributed](#) to deforestation of riparian corridors and human activities where there were once forests.

This is because riparian forests are the [link](#) between aquatic and [terrestrial ecosystems](#).

Riparian forests provide a wide range of ecosystem services and confer a great advantage on their adjacent surface freshwater. They [reduce](#)

climate change impacts by absorbing carbon dioxide, regulating the micro-climate and providing cover for [aquatic ecosystems](#). They also [stabilise](#) river banks, control erosion and siltation of inland waters, and provide food for [aquatic animals](#).

My research group recently [investigated](#) a Nigerian stream, Opa Stream, Ile-Ife, whose riparian corridor had been deforested. We found a worrying level of [heavy metal pollution](#) in the water. This is probably the result of human activities and related to the loss of forest.

Effects of deforestation

The deforestation of riparian corridor comes with two major impacts. First is the loss of [ecosystem services](#), which leads to the poor health of adjacent freshwater [ecosystems](#). Second is the negative [impact](#) on freshwater systems of human activities that take place on land converted from forest. Vegetation would mitigate heavy metal pollution because its presence means no agricultural or industrial activity on the riparian corridor. It also reduces the amount of run-off into freshwater systems, thus reducing the amount of human-generated pollutants.

A most noticeable impact is the release of persistent bio-accumulative toxicants to adjacent waters. These toxicants [include heavy metals](#) and organo-metal compounds, which accumulate in animal tissues at higher concentrations along the food chain.

My research group [found](#) five heavy metals—lead, cadmium, iron, arsenic and copper—exceeding their [freshwater recommended limits](#) in the stream we studied. We also found a strong linkage for concentrations of lead, copper and arsenic between the water column and the stream bed. This suggests that the entire freshwater system was contaminated with the metals and all the resident organisms were predisposed to contamination on account of the metals.

In this case, the heavy metal concentration is due to car-wash activities and block-making industries on the riparian corridor.

The use of pesticides on subsistence and commercial farms too contributed. Cadmium concentration was particularly high in water and sediment samples. High concentration of cadmium is another consequence of agricultural practice on the riparian corridor and it has varied level of [toxicity](#) among aquatic animals.

We do not have baseline data on this, but concentrations of these metals could have been heightened by prevalent activities on the riparian corridor.

We [reported](#) that a good number of the animals in the stream could be under stress because of the heavy [metal](#) pollution. These include the micro-organisms, plankton, insects and fishes, all of which can become less physiologically fit because of [heavy metal pollution](#). Plankton for instance [can take up heavy metals](#) in the water column, transfer such to bigger animals in the food chain where they become magnified in concentrations.

Restoring degraded tropical freshwater systems

It's important to conserve pristine or near-pristine tropical freshwater systems and restore the degraded ones. A good way to start would be by conserving or restoring the health of riparian forests, since this often [determines](#) the health of their adjacent water-bodies.

Riparian forests are an inseparable part of freshwater ecosystems, and should be protected by relevant government authorities at local, state and national levels.

People often think about what they can get from nature. They think of

food, water, shelter and economic gain—but they don't always think of the many ecosystem services they get that are less obvious. Human existence hinges on the visible and hidden ecosystem services of nature, including riparian [forest](#) and freshwater ecosystems.

This is why individuals, local communities and governments should do more to check the rate of human encroachment on riparian corridors. More effort is also needed to reclaim and reforest these areas.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Destroying Nigeria's riverside forests is bad for the freshwater ecosystem (2020, March 3) retrieved 27 April 2024 from <https://phys.org/news/2020-03-nigeria-riverside-forests-bad-freshwater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.