


Perovskite solution aging: Scientists find solution

March 17 2020



The clean stabilizer triethyl borate is used to inhibit side reactions in the perovskite solution. Credit: WANG Xiao

A research team from the Qingdao Institute of Bioenergy and Bioprocess Technology (QIBEBT) of the Chinese Academy of Sciences (CAS) has proposed a new understanding of the aging process of perovskite solution and also found a way to avoid side reactions.

Perovskite solar [cells](#) have developed quickly in the past decade. But like silicon solar cells, the efficiency of [perovskite solar cells](#) is highly dependent upon the quality of the [perovskite](#) layer, which is related to its crystallinity.

Unfortunately, the aging process of the perovskite [solution](#) used to fabricate solar cells makes the solution unstable, thus leading to poor efficiency and poor reproducibility of the devices. Reactants and preparation conditions also contribute to poor quality.

To combat these problems, a research team from the Qingdao Institute of Bioenergy and Bioprocess Technology (QIBEBT) of the Chinese Academy of Sciences (CAS) has proposed a new understanding of the aging process of perovskite solution and also found a way to avoid side reactions. The study was published in *Chem* on Mar. 17, titled "Perovskite solution aging: What happened and how to inhibit?"

Prof. PANG Shuping, corresponding author of the paper, said, "An in-depth understanding of fundamental solution chemistry" had not kept up with rapid efficiency improvements in perovskite solar cells, even though such cells have been studied for 10 years.

"Normally, we need high temperature and a long time to fully dissolve

the reactants, but some side reactions can happen simultaneously," said Prof. PANG. "Fortunately, we have found a way to inhibit them."

Achieving a highly stable perovskite solution is especially important in commercializing perovskite solar cells, since it will be easier to make devices with high consistency, said Prof. PANG.

WANG Xiao, an associate professor at QIBEBT and the first author of the paper, said side condensation reactions happen when methylammonium iodide and formamidinium iodide coexist in the solution. They represent the main side reactions in aging perovskite solution, although other side reactions between solute and solvent can occur at very high temperature.

FAN Yingping, a [graduate student](#) at the Qingdao University of Science and Technology (QUST) and the co-first author of the paper, studied many methods for stopping unwanted side reactions, but finally found that the stabilizer triethyl borate at a low boiling point was very effective. FAN also noted that it's a clean stabilizer, because it can be fully removed from the film during the following thermal annealing treatment.

With this new stabilizer, the reproducibility of the perovskite [solar cells](#) has improved greatly. "Now, we don't need to make fresh solutions every time before we make devices," said Prof. CUI Guanglei from QIBEBT, who noted that the finding is important for the fabrication of perovskite modules.

More information: Perovskite solution aging: What happened and how to inhibit? *Chem* (2020). [DOI: 10.1016/j.chempr.2020.02.016](https://doi.org/10.1016/j.chempr.2020.02.016)

Provided by Chinese Academy of Sciences

Citation: Perovskite solution aging: Scientists find solution (2020, March 17) retrieved 5 February 2023 from <https://phys.org/news/2020-03-perovskite-solution-aging-scientists.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.