

Study reveals a mechanism that plants can use to dissipate excess sunlight as heat

March 10 2020



Credit: CC0 Public Domain

For plants, sunlight can be a double-edged sword. They need it to drive photosynthesis, the process that allows them to store solar energy as sugar molecules, but too much sun can dehydrate and damage their

leaves.

A primary strategy that [plants](#) use to protect themselves from this kind of photodamage is to dissipate the extra light as heat. However, there has been much debate over the past several decades over how plants actually achieve this.

"During photosynthesis, light-harvesting complexes play two seemingly contradictory roles. They absorb [energy](#) to drive water-splitting and photosynthesis, but at the same time, when there's too much energy, they have to also be able to get rid of it," says Gabriela Schlau-Cohen, the Thomas D. and Virginia W. Cabot Career Development Assistant Professor of Chemistry at MIT.

In a new study, Schlau-Cohen and colleagues at MIT, the University of Pavia, and the University of Verona directly observed, for the first time, one of the possible mechanisms that have been proposed for how plants dissipate energy. The researchers used a highly sensitive type of spectroscopy to determine that [excess energy](#) is transferred from chlorophyll, the pigment that gives leaves their green color, to other pigments called carotenoids, which can then release the energy as heat.

"This is the first direct observation of chlorophyll-to-carotenoid energy transfer in the light-harvesting complex of green plants," says Schlau-Cohen, who is the senior author of the study. "That's the simplest proposal, but no one's been able to find this photophysical pathway until now."

MIT graduate student Minjung Son is the lead author of the study, which appears today in *Nature Communications*. Other authors are Samuel Gordon '18, Alberta Pinnola of the University of Pavia, in Italy, and Roberto Bassi of the University of Verona.

Excess energy

When sunlight strikes a plant, specialized proteins known as light-harvesting complexes absorb [light energy](#) in the form of photons, with the help of pigments such as chlorophyll. These photons drive the production of [sugar molecules](#), which store the energy for later use.

Much previous research has shown that plants are able to quickly adapt to changes in sunlight intensity. In very sunny conditions, they convert only about 30 percent of the available sunlight into sugar, while the rest is released as heat. If this excess energy is allowed to remain in the plant cells, it creates harmful molecules called free radicals that can damage proteins and other important cellular molecules.

"Plants can respond to fast changes in solar intensity by getting rid of extra energy, but what that photophysical pathway is has been debated for decades," Schlau-Cohen says.

The simplest hypothesis for how plants get rid of these extra photons is that once the light-harvesting complex absorbs them, chlorophylls pass them to nearby molecules called carotenoids. Carotenoids, which include lycopene and beta-carotene, are very good at getting rid of excess energy through rapid vibration. They are also skillful scavengers of [free radicals](#), which helps to prevent damage to cells.

A similar type of energy transfer has been observed in bacterial proteins that are related to chlorophyll, but until now, it had not been seen in plants. One reason why it has been hard to observe this phenomenon is that it occurs on a very fast time scale (femtoseconds, or quadrillionths of a second). Another obstacle is that the energy transfer spans a broad range of energy levels. Until recently, existing methods for observing this process could only measure a small swath of the spectrum of visible light.

In 2017, Schlau-Cohen's lab developed a modification to a femtosecond spectroscopic technique that allows them to look at a broader range of energy levels, spanning red to blue light. This meant that they could monitor energy transfer between chlorophylls, which absorb red light, and carotenoids, which absorb blue and green light.

In this study, the researchers used this technique to show that photons move from an excited state, which is spread over multiple chlorophyll molecules within a light-harvesting complex, to nearby carotenoid molecules within the complex.

"By broadening the spectral bandwidth, we could look at the connection between the blue and the red ranges, allowing us to map out the changes in energy level. You can see energy moving from one [excited state](#) to another," Schlau-Cohen says.

Once the carotenoids accept the excess energy, they release most of it as heat, preventing light-induced damage to the cells.

Boosting crop yields

The researchers performed their experiments in two different environments—one in which the proteins were in a detergent solution, and one in which they were embedded in a special type of self-assembling membrane called a nanodisc. They found that the energy transfer occurred more rapidly in the nanodisc, suggesting that environmental conditions affect the rate of energy dissipation.

It remains a mystery exactly how excess sunlight triggers this mechanism within plant cells. Schlau-Cohen's lab is now exploring whether the organization of chlorophylls and carotenoids within the chloroplast membrane play a role in activating the photoprotection system.

A better understanding of plants' natural photoprotection system could help scientists develop new ways to improve crop yields, Schlau-Cohen says. A 2016 paper from University of Illinois researchers showed that by overproducing all of the proteins involved in photoprotection, [crop yields](#) could be boosted by 15 to 20 percent. That paper also suggested that production could be further increased to a theoretical maximum of about 30 percent.

"If we understand the mechanism, instead of just upregulating everything and getting 15 to 20 percent, we could really optimize the system and get to that theoretical maximum of 30 percent," Schlau-Cohen says.

Provided by Massachusetts Institute of Technology

Citation: Study reveals a mechanism that plants can use to dissipate excess sunlight as heat (2020, March 10) retrieved 23 July 2024 from <https://phys.org/news/2020-03-reveals-mechanism-dissipate-excess-sunlight.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.