


# Valley reshaping and damming change surface and groundwater nitrate level at watershed scale

March 10 2020, by Li Yuan



Land use and land cover types in the valley of the treated watershed before and after the project of Gully Land Consolidation (GLC) project. The remote sensing images show that the valley flat farmland significantly expanded, and reservoirs were built in the creek bottom. Credit: YU Yunlong, et al.

On the Chinese Loess Plateau (CLP), a project named "Gully Land Consolidation" (GLC) has been implemented in rural watersheds to offset the farmland loss induced by "Grain for Green Program". Its main

goal includes reshaping creek valleys by incising foot slopes, filling stream channels and ditches, constructing drainage canals, dams and reservoirs, and creating flat farmlands.

Previous studies found that, compared to a natural catchment, the [soil nitrate](#) increased significantly in the deep profile of the [valley](#) flat farmland in a reshaped catchment. However, whether such valley reshaping, damming and agricultural land consolidation affect the [surface](#) and [groundwater nitrate](#) and pose a threat to local people remains unclear.

Recently, Prof. Jin Zhao's team from the Institute of Earth Environment (IEE) of the Chinese Academy of Sciences conducted extensive field investigation and experiments to enhance the understanding of water chemistry, inorganic nitrogen and environmental isotope levels in a dammed [watershed](#) and a natural watershed, revealing how valley shaping and damming affect water nitrate level and source at watershed scale.

They found that nitrate concentrations in both the natural and dammed watersheds were lower than  $10 \text{ mg}\cdot\text{L}^{-1}$ , among which the seasonal nitrate contents in the surface ( $0.42\text{-}0.52 \text{ mg}\cdot\text{L}^{-1}$ ) and groundwater ( $1.86\text{-}3.65 \text{ mg}\cdot\text{L}^{-1}$ ) of the dammed watershed were significantly lower than those of the natural watershed (surface water:  $0.92\text{-}2.07 \text{ mg}\cdot\text{L}^{-1}$ ; groundwater:  $2.73\text{-}8.21 \text{ mg}\cdot\text{L}^{-1}$ ).

Hierarchical cluster analysis (HCA) showed that there were strong interactions between surface runoff and the reservoir water in the dammed watershed, while in the natural watershed, streamflow had a close relationship with groundwater.

They also found that nitrate fertilizer, soil ammonium, and manure and septic waste were the main sources of nitrate in the surface water of two

watersheds, while the nitrate in the groundwater mainly originated from the nitrification of soil ammonium and manure and septic waste.

Compared to the natural watershed, the dammed watershed had increased inputs of agricultural fertilizers in the surface water. However, there was no significant difference in the groundwater nitrate sources between the two watersheds.

They considered that valley reshaping, damming and agricultural land consolidation significantly affected the input of each nitrate source and water quality of the [surface water](#) but had little influence on the groundwater.

Moreover, the study revealed that valley reshaping, damming and agricultural land consolidation could alleviate nitrate pollution and that dams and reservoirs built along the reshaped valley would play a key role in controlling the concentrations of water nitrate, which had global implications for watershed management and agricultural land consolidation.

However, although the current groundwater had not been influenced noticeably by GLC projects in the dammed watershed, the time lag between agricultural land consolidation and groundwater quality response should be considered in the future.

"A reasonable level of N application should be implemented in the dammed watershed to maintain the high-quality of the surface and groundwater," said Prof. Jin.

This study aids in developing strategies for the comprehensive management of watershed and drinking water sources in rural regions worldwide.

**More information:** Yunlong Yu et al. Effects of valley reshaping and damming on surface and groundwater nitrate on the Chinese Loess Plateau, *Journal of Hydrology* (2020). [DOI: 10.1016/j.jhydrol.2020.124702](https://doi.org/10.1016/j.jhydrol.2020.124702)

Provided by Chinese Academy of Sciences

Citation: Valley reshaping and damming change surface and groundwater nitrate level at watershed scale (2020, March 10) retrieved 20 April 2024 from <https://phys.org/news/2020-03-valley-reshaping-surface-groundwater-nitrate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.