


Belle II yields the first results: In search of the Z' boson

April 7 2020



A computer graphics of a simulated event in which a Z' boson is produced by e^+e^- collisions, in association with two muons (green line and hits) and decays into invisible particles. In this figure, the Z' boson decays into an invisible neutrino and anti-neutrino, but it may decays also into the dark matter particle and its anti-particle. Credit: KEK / Belle II Collaboration

The Belle II experiment has been collecting data from physical measurements for about one year. After several years of rebuilding work, both the SuperKEKB electron–positron accelerator and the Belle II detector have been improved compared with their predecessors in

order to achieve a 40-fold higher data rate.

Scientists at 12 institutes in Germany are involved in constructing and operating the detector, developing evaluation algorithms and analyzing the data. The Max Planck Institute for Physics made a substantial contribution to the development of the highly sensitive innermost detector, the Pixel Vertex Detector.


With the help of Belle II, scientists are looking for traces of new physics that could explain the unequal occurrence of matter and antimatter and the mysterious [dark matter](#). One of the so-far undiscovered particles that the Belle II detector is looking for is the Z' boson—a variant of the Z boson, which acts as an exchange particle for the weak interaction.

As far as we know, about 25% of the universe consists of dark matter, whereas visible matter accounts for just under 5% of the energy budget. Both forms of matter attract each other through gravity. Dark matter thus forms a kind of template for the distribution of visible matter. This can be seen, for example, in the arrangement of galaxies in the universe.

Link between dark and normal matter

The Z' boson may play an interesting role in the interaction between dark and visible matter, (i.e., it could be a kind of mediator between the two forms of matter). The Z' boson can—at least theoretically—result from the collision of electrons (matter) and positrons (anti-matter) in the SuperKEKB and then decay into invisible dark matter particles.

The Z' boson can thus help scientists to understand the behavior of dark matter. What's more, the discovery of the Z' boson could also explain other observations that are not consistent with the [standard model](#), the fundamental theory of particle physics.



Electrons and positrons collide within the Belle II detector. Credit: ill./©: Belle II

Important clue: Detection of muon pairs

But how can the Z' boson be detected in the Belle II detector? Not directly—that much is sure. Theoretical models and simulations predict that the Z' boson could reveal itself through interactions with muons, the heavier relatives of electrons. If scientists discover an unusually high number of muon pairs of opposite charge after the electron/positron collisions as well as unexpected deviations in energy and momentum conservation, this would be an important indication of the Z' boson.

However, the new Belle II data has not yet provided any indication of the Z' boson. But with the new data, the scientists can limit the mass and coupling strengths of the Z' [boson](#) with previously unattainable accuracy.

More data, more precise analyses

"Despite the still small amount of data, we can now make measurements that have never been done before," says the spokesperson of the German groups, Dr. Thomas Kuhr from the Ludwig Maximilian University of Munich. "This underlines the important role of the Belle II experiment in the study of elementary particles."

These initial results come from the analysis of a small amount of data collected during the start-up phase of SuperKEKB in 2018. Belle II went into full operation on March 25, 2019. Since then, the experiment has been collecting data while continuously improving the collision rate of electrons and positrons.

If the experiment is perfectly tuned, it will provide considerably more data than in the recently published analyses. The physicists thus hope to gain new insights into the nature of dark matter and other unanswered questions.

More information: I. Adachi et al. Search for an Invisibly Decaying Z' Boson at Belle II in $e+e\rightarrow\mu+\mu-(e\pm\mu\mp)$ Plus Missing Energy Final States, *Physical Review Letters* (2020). [DOI: 10.1103/PhysRevLett.124.141801](https://doi.org/10.1103/PhysRevLett.124.141801)

Provided by Max Planck Society

Citation: Belle II yields the first results: In search of the Z' boson (2020, April 7) retrieved 23 March 2023 from <https://phys.org/news/2020-04-belle-ii-yields-results-boson.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.