

# Bermudagrass harvest management options with poultry litter fertilization

April 30 2020

---



Regrowth of two hybrid bermudagrass [*Cynodon dactylon* (L.) Pers.] varieties after cutting at different time intervals and stubble heights. Credit: John Read

When fertilizing bermudagrass with poultry litter, turfgrass managers must consider limiting the buildup of soil P or drawing down soil test P through cut-and-carry forage. In a previous study that provided turfgrass with 122 kg ha<sup>-1</sup> P in poultry litter, researchers found that Tifton 44 bermudagrass cut every 49 days at 3-cm stubble height recovered 23% of the P applied. Bermudagrass P removal is closely associated with dry matter (DM) yield and can be altered through management variables like variety, plant maturity at harvest, and litter application rate or timing. Given the tradeoff between DM yield and the amounts of crude protein

and digestible DM in bermudagrass, managing harvests to maximize P recovery may adversely impact forage nutritive value.

In an article recently published in *Crop, Forage & Turfgrass Management*, researchers report on the response of [forage](#) nutritive value and P removal to the combined effects of [harvest](#) interval and stubble height in Russell and Tifton 44 bermudagrass receiving 4 tons acre<sup>-1</sup> of [poultry litter](#) supplemented with 98 lb acre<sup>-1</sup> N each spring from 2005 to 2007. Soil test P increased by approximately 60 mg kg<sup>-1</sup>, but there was no difference in soil P accumulation in the harvest interval x stubble height plots in autumn 2007. They concluded a best management practice is cutting every 35 d at 3-cm residual stubble height, which has long been considered a standard for economical forage production. Additionally, harvesting bermudagrass at an advanced stage of maturity is a best management practice in situations where high soil test P is of greater concern than forage nutritive value.

**More information:** John J. Read et al, Managing harvest of 'Russell' and 'Tifton 44' bermudagrass receiving broiler litter for nutritive value and phosphorus removal, *Crop, Forage & Turfgrass Management* (2020). [DOI: 10.1002/cft2.20013](https://doi.org/10.1002/cft2.20013)

Provided by American Society of Agronomy

Citation: Bermudagrass harvest management options with poultry litter fertilization (2020, April 30) retrieved 27 June 2024 from <https://phys.org/news/2020-04-bermudagrass-harvest-options-poultry-litter.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|