

Researchers provide new clue to solar coronal heating problem

April 6 2020, by Liu Jia



Credit: CC0 Public Domain

Coronal heating is a topic dedicated to explaining how the corona may be heated up to a temperature of millions of degrees, far above that of the photosphere. To transport magnetic energy into the corona, Alfvén-



wave turbulence is a promising candidate. However, how the wave is generated in the corona is still an open question.

In a study published online in the *Astrophysical Journal Letters*, Dr. Bi Yi and his collaborators from Yunnan observatories of the Chinese Academy of Sciences revealed that Alfvénic wave energy communicated by the descending knots in the solar prominence may well contribute in a non-negligible way to the local heating of the surrounding <u>corona</u>.

This investigation is based on the observation from NVST, which is a ground-based multichannel high-resolution imaging system, including the H α , G-band, and TiO-band. The NVST H α data is useful to study the dynamics of quiescent prominences. The investigated prominence showed that the descending and rising knots in the prominence investigated are colored.

The researchers also tried to estimate the loss rate of the gravitational potential from the descending knots and compared it with heating power requirement for the corona in the quiet Sun. After investigating the size, velocity and path of thousands of the falling knots in this prominence, they found that the gravitational energy loss rate of these observed knots amounts to about 1/2000 of that required to heat the entire quiet Sun, increasing to 1/320 when considering possibly further downward motions of the knots having disappeared in the H α observations.

Assuming that the descending knots were capable of exciting Alfvén waves that could then dissipate within the local corona, the gravitational potential energy of the knots may have been converted into thermal energy. Such a mechanism may contribute to the heating of the corona local to these prominences.

A robust estimation of the density of the knots in the prominence will enable the precise estimation of the loss rate of the gravitational



potential of the falling plasma. Additional modeling work is also required in order to reveal how the interaction between the falling dense plasma and the <u>magnetic field</u> generates the Alfvén wave, dissipation of which is capable of contributing to the heating of the local corona.

More information: Yi Bi et al. Dynamics of Descending Knots in a Solar Prominence and Their Possible Contributions to the Heating of the Local Corona, *The Astrophysical Journal* (2020). DOI: 10.3847/2041-8213/ab79a2

Provided by Chinese Academy of Sciences

Citation: Researchers provide new clue to solar coronal heating problem (2020, April 6) retrieved 2 May 2024 from <u>https://phys.org/news/2020-04-clue-solar-coronal-problem.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.