


## Researchers watch the dynamics of plasmonic skyrmions made from light on ultra-smooth gold platelets for the first time

April 24 2020



Snapshot of the nanofilm that displays the electric field of light in the plasmonic nanovortices (skyrmions). The hexagonal (6-fold) symmetry in the plane is easily



visible. Credit: University of Stuttgart

The destructive force of a tornado occurs due to the extremely high rotational speeds in its center, which is called a vortex. Surprisingly, similar effects are predicted for light that travels along an atomically smooth gold surface, which can exhibit angular momentum and vortices. Researchers at the Universities of Stuttgart and Duisburg-Essen and the University of Melbourne (Australia) have now succeeded for the first time in filming these vortex patterns, which are called skyrmions, on the nanometer scale. The journal *Science* reports this groundbreaking work in its issue of April 24, 2020.


When a figure skater starts to perform a pirouette and raises her arms, she turns around her own axis increasingly faster due to the conservation of angular momentum. On hot summer days, the same pirouette effect creates so-called "dust devils," i.e., small whirlwinds of hot air, and it also gives large tornadoes their destructive power. The physicist Tony Skyrme studied such vortices in detail in the 1960s in a field of research called topology. The patterns are called skyrmions after their discoverer.

Light that travels along atomically smooth, nanostructured gold surfaces can also have a kind of angular momentum, and therefore vortices can form. However, in this case, the vortices are only a few hundred nanometers in size, and the eye of these nanostorm is only a few nanometers in size. Therefore, no one has yet been able to measure the exact orientation of these vortices. It was also impossible to look at the vortex dynamics, because the time it takes <u>light</u> to travel once around such a vortex is only a few femtoseconds (trillionths of a millisecond).

In a breakthrough experiment, a team of researchers from the University of Stuttgart, the University of Duisburg-Essen, and the University of



Melbourne in Australia has succeeded for the first time in filming such plasmonic skyrmions made of light on the nanometer scale. The researchers were able to record the direction of the electric and magnetic field of the light in all three dimensions, and even measured its dynamics. Theorist Tim Davis from Melbourne, who visited Stuttgart and Duisburg with support of the IQST quantum center, calculated the required light wavelengths, the optimal shape of the nanostructures, as well as the exact thickness of the gold platelets. He predicted how regular arrays (called <a href="mailto:skyrmion">skyrmion</a> lattices) of light vortices would behave.



Analogy to nano tornadoes in plasmonic vortices (skyrmions). The display exibits the measured 3-dimensional structure of the magnetic field of the light on the gold nanoplatelets. Credit: University of Stuttgart



Bettina Frank from the research group of Harald Giessen at the 4th Physics Institute at the University of Stuttgart produced atomically smooth gold platelets with adjustable thickness in the nanometer range using a newly developed method. Extremely flat silicon wafers were used as a substrate for this purpose. The gold platelets were nanostructured with a high-precision gold ion beam. When irradiated with laser pulses of a carefully calculated wavelength in the infrared range, entire skyrmion arrays of light, so-called plasmonic skyrmions, could be created.

The measurement of the vector dynamics, i.e., the three-dimensional alignment of the plasmon light fields and their temporal behavior, was successfully carried out in a specifically designed new experiment in the group of Frank Meyer zu Heringdorf at the University of Duisburg-Essen. Doctoral students Pascal Dreher and David Janoschka sent laser pulses of only 13 femtoseconds duration at 800-nm wavelength onto the gold platelets with the nanostructures. The photoeffect, for which Einstein received his Nobel Prize, causes electrons to be ejected out of the gold sample, which are then measured with an electron microscope. By smartly combining several laser pulses with different light polarizations and repeating the experiment several times, the vector components of the light fields could be determined by projection.

By sending two laser pulses one after another onto the sample, the nanotornadoes of light can both be excited and subsequently probed by ultrashort <u>laser pulses</u>, so that within the duration of about one night, an entire nanofilm of these light vortices can be recorded.

Harald Giessen from Stuttgart believes that it might be possible in the future on the basis of this research to create new types of microscopes that could produce much smaller structures with light than has been the case so far. "The combination of the orbital angular momentum and the vector properties leads to plasmonic vortex structures in the nanometer



range even with linear optics," he reports. "It will also be possible to experimentally observe time-resolved skyrmion physics under a variety of boundary conditions."

The interaction of such skyrmion fields and their orbital <u>angular</u> <u>momentum</u> with neighboring particles in semiconductors, for example in atomically thin, two-dimensional materials, will be particularly exciting. "Thanks to our new Raith ion-beam lithography machine, we have almost infinite possibilities for generating different topological nanostructures and studying their skyrmion dynamics with the Duisburg nanocamera."

**More information:** Ultrafast vector imaging of plasmonic skyrmion dynamics with deep subwavelength resolution. *Science* 24 Apr 2020: DOI: 10.1126/science.aba6415

## Provided by University of Stuttgart

Citation: Researchers watch the dynamics of plasmonic skyrmions made from light on ultrasmooth gold platelets for the first time (2020, April 24) retrieved 24 April 2024 from <a href="https://phys.org/news/2020-04-dynamics-plasmonic-skyrmions-ultra-smooth-gold.html">https://phys.org/news/2020-04-dynamics-plasmonic-skyrmions-ultra-smooth-gold.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.