

Reference genes identified that are useful for genetic improvement in wheat

April 21 2020



Credit: CC0 Public Domain

Wheat feeds the world. According to the FAO, wheat is one of the world's main crops, both in terms of extent and production, as well as being one of the main sources of carbohydrates and vegetable protein in

the human diet. The quest for genetic improvement in wheat, leading to varieties that are more resistant to issues brought about by climate change or certain pests, is a response to the need to keep feeding people.

Genetic improvement programs can span decades and entail studying genetic mechanisms of wheat in order to better understand them and hence accomplish more effective research in this field. In this context, a research team made up of Professor Miguel Aguilar, from the Department of Botany, Ecology and Plant Physiology at the University of Cordoba, along with José Garrido and Pilar Prieto, both researchers at the Institute for Sustainable Agriculture (abbreviated to IAS—CSIC in Spanish), was able to further our knowledge of these procedures. They published the set of reliable reference genes in wheat meiosis.

Meiosis is the division process that generates gametes or reproductive cells. During meiosis, chromosome association and genetic recombination occur. These are vital processes for a plant's fertility. Reference genes or normalizing genes are genes whose expression does not change during the biological processes under study. So, these kinds of genes work as a constant in order to ensure that any change of expression observed in other genes can be significantly correlated with the studied process.

The absence of validated reference genes that can be used to study wheat meiosis is what led this team to seek the set of genes that can provide precise measurements of the expression of other genes during the process of meiosis in [durum wheat](#), used to make pasta, and common wheat, used to make bread. This discovery will be useful for future scientific research focused on the [genetic improvement](#) of wheat.

Using the recently sequenced and assembled complete wheat genome as a basis, the research team used the quantitative PCR technique (used to amplify DNA sequences) to test and analyze the stability of certain genes

and likewise determine how many are needed for the results to be reliable. The reference genes obtained were duplicated or were part of a gene family, so the real challenge for the team was to identify the specific members that can be used as reference genes in gene expression during meiosis. Lastly, two new genes were identified: the cyclic phosphodiesterase-like gene and the salt tolerant protein gene.

This set of genes helps us to understand how meiosis works and how we can manipulate it, which could revolutionize agriculture, since it would allow for manipulating chromosome associations and using [genes](#) from other species in [wheat](#)—all this to continue the job of feeding the world.

More information: José Garrido et al, Identification and validation of reference genes for RT-qPCR normalization in wheat meiosis, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-59580-5](https://doi.org/10.1038/s41598-020-59580-5)

Provided by University of Córdoba

Citation: Reference genes identified that are useful for genetic improvement in wheat (2020, April 21) retrieved 25 April 2024 from <https://phys.org/news/2020-04-genes-genetic-wheat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.