

# Grassland studies, radar-tracked bumblebees offer clues for protecting pollinators

April 27 2020, by Anthony King, From Horizon Magazine

---



It's not just crops that rely on bumblebees; wild plants do too. Credit: Anthony King

Scientists used a radar to track a bumblebee from its maiden flight until death for the first time as part of wider research racing to understand the

impact and needs of declining bee populations, including on Europe's fragmented biodiversity hotspots—grasslands.

Wild bees are major pollinators of crops in Europe such as [peas, beans and oilseed rape](#). But as [their numbers decline](#), there is a need to monitor and boost [bee populations](#) and understand the consequences of fewer bees.

And pollinators are not just important for crops—but also [wild plants](#). [Falling bee numbers](#) are likely already having an impact—we just don't know exactly what that is.

"We have seen declines in pollinators as [land use](#) has intensified," said Professor Yann Clough, ecologist at Lund University in Sweden. "It is not just crops, but wild plants which they pollinate, which begs the question, do declines in pollinators mean we will see declines in the reproduction of wild plants?"

Some plants such as grasses can thrive without pollinating insects because they are wind pollinated. Others rely exclusively on insects to reproduce. Many wildflowers especially are pollinated by [wild bees](#) such as bumblebees. This includes vetches, peas such as yellow bird's-foot trefoil and many types of daisy.

"About one-third of plants do not at all depend on insects for pollination, and a third do, to some extent," Prof. Clough said. "Another third then are highly dependent on insects for pollination."

Without bees, ecosystems could tilt towards plants that do not need bees, nor cater for them. Fewer bees would mean less diversity in wild plants and pollinators and could impact birds and mammals that depend on them.

Prof Clough is studying two dozen flower-rich grasslands in Sweden to better understand how the landscapes surrounding these sites influence bee numbers and the plants growing there through a project called [DrivenByPollinators](#).



Europe's fragmented grasslands are as diverse as rainforests, says Prof. Yann Clough, an ecologist at Sweden's Lund University. Credit: Yann Clough

## Grasslands

The semi-natural grasslands Prof. Clough studies in Sweden are

'unimproved,' meaning they were probably never ploughed or fertilised. "They tend to have boulders in them," he said. Cattle, sheep and goats graze there.

Over the past 150 years, semi-natural grasslands in Sweden have reduced by about 90%, he says. They are now mostly small fragments, oases of biodiversity in a sea of intensively managed fields of crops such as potato or wheat.

"These semi-natural grasslands are like our rainforests, in terms of the density of species and the rarity of some species here," said Prof. Clough. This is because a diverse community of flowers, insects, birds and other animals live and depend on one another in these habitats.

It is likely that what happens in these grasslands will depend strongly on the surrounding countryside, since insect pollinators are mobile and many bees rely on pollen and nectar in the wider landscape.

"We are concerned that the quality of these fragments might decrease due to the indirect effects of intensive management in the surrounding landscape," said Prof. Clough.

To study the impact of the countryside, he and his team will measure over several years the seeds produced and number of seedlings in the grasslands. In one experiment, Prof. Clough is shutting out pollinators from small sections of the wild fields with a tent-like netting-covered dome structure to study the impact on plant communities.

A fall in wild bees and other pollinators could help change [grassland](#) composition, for example. "That might lead to a shift towards more grasses, and less legumes, which fix nitrogen in these grasslands," said Prof. Clough. Legumes plants such as peas and clover take unreactive nitrogen from the air, and with the assistance of microbial partners on

their roots, turn it into nitrogen-rich fertiliser compounds for other plants.

This could have knock-on effects for the ecosystem and for farmers. More grasses and fewer flowers mean drier and less nutritious fodder for grazing animals.

Farmers can receive subsidies for grazing grasslands to conserve biodiversity. If, within a few years, agriculture policy in Europe moves towards a more [results-based payment](#) for biodiversity friendly management, farmers will want to be sure that the biodiversity in their grasslands is not slowly eroding because of the surrounding landscape, says Dr. Clough.

## **Flower strips**

Prof. Clough hopes to use his results to draw up recommendations on how to best manage the landscapes around grasslands for pollinators such as bumblebees. "It might mean conserving more natural habitat around some grasslands, or putting in (pollinator friendly) flower strips in the wider landscape, or creating (bee) nesting sites by planting hedge or wood habitats," he said.

But studying exactly how declining pollinator populations affect biodiversity is a challenge, as is studying any changes to bee behaviour and foraging.

One way to study an animal's behaviour is to track it. For large mammals such as bears or deer, a collar can be placed around their neck to follow their movements. But with bumblebees, this is tricky.

At an agriculture research station north of London, however, scientists were able to glue a lightweight harmonic radar to the backs of bees. It

acts as a signal deflector which allows the bee to be traced as they forage. A radar dish emits a signal, which bounces off the transponder, and this signal is picked up by a receiving radar. The position of the bee is pinpointed every few seconds.

"With two different radars mounted on the same (truck) trailer, that allows us track the bees up to a kilometre away," said [Professor Lars Chittka](#), behavioural ecologist at Queen Mary University of London, UK, and leader of the [SpaceRadarPollinator](#) project, which tracked the bees.

The team aims now to build multiple such radars and place them over a larger area to track bees for longer distances.

## **Altitude**

Until now, they have only tracked the position but not the altitude of the bee. "The biggest breakthrough is the [3-dimensional tracking](#)," said Prof. Chittka, which he hopes to test in the summer. "That will allow us reconstruct what the bee sees while in flight. We can see how it memorises landmarks and uses them (for orientation)."

## **A 3-D video simulation of a bee out to forage**

Bee tracking could allow scientists to offer advice on how best to conserve pollinators. "It is important for us to understand bees' foraging needs," said Prof. Chittka. "Not just what resources they require, but also how they should be distributed in space."

It might be better to have strips of connected habitat planted along field margins or the edges of roads, for example.

"We hope to get exact predictions of how far apart habitats are allowed

to be, or at what point foraging behaviour or dispersal of queens might be disrupted," he said.

The project for the first time ever tracked a bumblebee from maiden flight until death. Bees initially explored their surroundings, departing on orientation flights in different directions and looping back to their nest. Typically, they then located a profitable flower patch and spent lots of time revisiting it.

But not all bees follow the same strategy. "Some individuals were relatively faithful to flower patches, while others were wanderers all their lifetime," said Prof. Chittka.



For the first time ever, researchers tracked a bumblebee from its maiden flight to

death. Credit: Yann Cloug

Four bumblebees were tracked as part of the project. Most lived three to four weeks, before disappearing off the radar, presumed eaten by birds. "We give them a number, like 007, to identify them," Prof. Chittka said.

Prof. Clough says that little is still known about how pollinators use the landscape, which makes it difficult to know what pollinator-friendly steps we can take.

He says that technical developments that shed more light into their movements are very much welcome. "The generated knowledge could be used to inform models for pollinator foraging, such as those that we are using in our research," he said.

Provided by Horizon: The EU Research & Innovation Magazine

Citation: Grassland studies, radar-tracked bumblebees offer clues for protecting pollinators (2020, April 27) retrieved 8 May 2024 from <https://phys.org/news/2020-04-grassland-radar-tracked-bumblebees-clues-pollinators.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.