

Harnessing psyllid peptides to fight citrus greening disease

April 28 2020



Asian citrus psyllids feed on a citrus tree. The psyllids deposit a bacterium in the sap that causes citrus greening disease, a scourge to the citrus industries in Florida and California, worth a combined \$17 billion. Credit: BTI

Citrus greening disease, also called huanglongbing (HLB), is a bacterial



infection of citrus trees that results in small, misshapen and sour fruits that are unsuitable for consumption, ultimately killing the tree. Because there is no cure, HLB is a major threat to the \$10 billion citrus industry in Florida, where it was first detected in 2005, and to the \$7 billion industry in California, where it appeared last year.

Researchers from the Boyce Thompson Institute, the U.S. Department of Agriculture Agricultural Research Service (USDA-ARS) and the University of Washington (UW), led by Michelle Heck, investigated a seemingly unlikely source of biocontrols for HLB: neuropeptides found in the insect that carries the disease-causing bacterium. Asian citrus psyllids (Diaphorina citri) infect <u>citrus trees</u> with Candidatus Liberibacter asiaticus (CLas) bacterium by feeding on their leaves and stems.

Laura Fleites, a USDA T.W. Edminster Research Associate in Heck's group at BTI, focused on neuropeptides because they function as hormones in hemipteran insects—a class that includes psyllids, aphids, whiteflies, shield bugs and other crop-plaguing species—to regulate growth, development and other biological functions. Moreover, other studies have shown that analogs of insect-derived neuropeptides kill pea aphids, and insecticides based on the kinin family of neuropeptides have been developed.

"If we could develop an insecticide that is specific for Asian citrus psyllids based on one of the insect's own neuropeptides, then we could protect citrus trees from the insect that spreads CLas," Heck said. "Citrus greening disease is devastating our <u>citrus industry</u>, and we need to develop new ways for our citrus growers to control it."

As reported in the April 3 issue of *Journal of Proteome Research*, Fleites and the team characterized the full array of peptides found in the psyllids and identified 122 candidate neuropeptides. While promising,



the findings offer only potential starting points for combatting HLB, because unmodified insect-derived neuropeptides lack the stability, bioavailability and half-lives needed for use as insecticides in the citrus grove.

To turn the findings into a usable product, the team is now collaborating with Ronald Nachman, a USDA researcher at College Station, Texas, Robert Shatters, a USDA researcher in Fort Pierce Florida, and Mark Trimmer from AgroSource, Inc, a small agritech company, to identify the best psyllid-derived neuropeptide for development. Then, the team will stabilize the peptide and decide the optimum method for delivering the insecticidal molecule to citrus trees—whether as a spray, by engineering trees to make the peptides themselves, or some other method.

For the study, Fleites, developed new extraction and analysis methods that other researchers could use in their investigations of insect peptides.

"Thanks to USDA support, I was able to develop techniques that enable the identification of small, functional insect peptides separately from their larger, inactive precursors," said Fleites. "Because these techniques aren't specific to psyllids, they may be useful for identifying neuropeptides in other hemipteran insects to protect crop plants."

Heck noted that this work shows how USDA grant programs can be used to fund everything from discovery to product development. "We performed the <u>neuropeptide</u> discovery in a study funded by NIFA, and are now doing the translational work under a different USDA grant program from APHIS," she said.

More information: Laura A. Fleites et al, Peptidomics Approaches for the Identification of Bioactive Molecules from Diaphorina citri, *Journal of Proteome Research* (2020). <u>DOI:</u>



10.1021/acs.jproteome.9b00509

Provided by Boyce Thompson Institute

Citation: Harnessing psyllid peptides to fight citrus greening disease (2020, April 28) retrieved 21 May 2024 from <u>https://phys.org/news/2020-04-harnessing-psyllid-peptides-citrus-greening.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.