


Major outburst and X-ray pulsations detected from RX J0209.6–7427

April 15 2020, by Tomasz Nowakowski



X-ray lightcurve of RX J0209.6–7427 based on NICER count rates (0.5–8.0 keV) averaged over 6 hour intervals. Credit: Vasilopoulos et al., 2020.

An international team of astronomers has identified a major outburst and X-ray pulsation from an X-ray binary system known as RX

J0209.6–7427. The detection, detailed in a paper published April 6 on the arXiv pre-print server, could shed more light on the nature of this source.

Generally, X-ray binaries are composed of a normal star or a white dwarf transferring mass onto a compact neutron star or a black hole. Based on the mass of the companion star, astronomers divide them into low-mass X-ray binaries (LMXB) and high-mass X-ray binaries (HMXB).

Of special interest are Be/X-ray binaries (BeXRBs), a subclass of HMXBs in which the optical star is a dwarf, subgiant or giant OBe star. Studying X-ray outbursts from BeXRBs could be essential in order to improve our understanding of the nature of X-ray binaries and their behavior.

Located in the outer wing of the Small Magellanic Cloud (SMC), RX J0209.6–7427 is a BeXRB first detected by the ROSAT satellite. The source exhibited two luminous outbursts, in March and November 1993, with measured luminosity at a level of about 100 undecillion erg/s.

Now, a group of astronomers led by Georgios Vasilopoulos of Yale University reports the detection of another significant outburst from RX J0209.6–7427 using the Monitor of All-sky X-ray Image (MAXI) instrument on the International Space Station (ISS). Moreover, the researchers conducted follow-up observations of this outburst with the Neutron Star Interior Composition Explorer (NICER) on ISS, the Nuclear Spectroscopic Telescope Array (NuSTAR) and Fermi spacecraft, which led to the discovery of X-ray pulsations from the system.

"In this paper, we present the first results of the X-ray monitoring of the system during its 2019 outburst. (...) We present the timing and spectral

analysis that resulted in the discovery of coherent pulsations and characterization of its broadband X-ray spectrum from NuSTAR and NICER data," the paper reads.

The outburst started on November 20, 2019, when when MAXI identified an uncatalogued X-ray transient source. Afterward, follow-up observations of this event commenced using NICER, NuSTAR and Fermi.

During the November 2019 outburst, the maximum X-ray luminosity of RX J0209.6–7427 reached 2.0 duodecillion erg/s, and NICER detected X-ray pulsations with a period of 9.3 seconds. Therefore, the system temporarily became a pulsating ultra-luminous X-ray source (PULX).

The high luminosity of RX J0209.6–7427 measured during the latest [outburst](#) suggests that it might be the brightest BeXRB ever observed in the SMC.

However, the astronomers noted that despite the super-Eddington luminosity of RX J0209.6–7427, the neutron star in the system appears to have only a moderate magnetic field strength. The researchers found that the surface magnetic field of the neutron star is at a level of approximately 3.0 trillion G and its orbital period is likely greater than 50 days.

More information: The 2019 super-Eddington outburst of RX J0209.6-7427: Detection of pulsations and constraints on the magnetic field strength, arXiv:2004.03022 [astro-ph.HE]
arxiv.org/abs/2004.03022

Citation: Major outburst and X-ray pulsations detected from RX J0209.6–7427 (2020, April 15)
retrieved 26 April 2024 from

<https://phys.org/news/2020-04-major-outburst-x-ray-pulsations-rx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.