


# Oysters and clams can be farmed together

April 2 2020



The four bivalve species studied at Rutgers' New Jersey Aquaculture Innovation Center include the Eastern oyster, Atlantic surfclam, hard clam and softshell clam (left to right). Credit: Michael Acquafredda/Rutgers University

Eastern oysters and three species of clams can be farmed together and

flourish, potentially boosting profits of shellfish growers, according to a Rutgers University-New Brunswick study.

Though diverse groups of [species](#) often outperform single-species groups, most bivalve farms in the United States and around the world grow their crops as monocultures, notes the study in the journal *Marine Ecology Progress Series*.

"Farming multiple species together can sustain the economic viability of farm operations and increase profitability by allowing shellfish growers to more easily navigate market forces if the price of each individual crop fluctuates," said lead author Michael P. Acquafredda, a doctoral student based at Rutgers' Haskin Shellfish Research Laboratory in Port Norris, New Jersey.

Farming mollusks such as clams, oysters and scallops contributes billions of dollars annually to the world's economy. In the United States, more than 47 million pounds of farm-raised clam, oyster and mussel meat worth more than \$340 million were harvested in 2016, the study says.

The study, which took place in a laboratory setting at Rutgers' New Jersey Aquaculture Innovation Center in North Cape May, New Jersey, tested the feasibility of farming multiple bivalve species in close proximity to each another.

Mimicking farm conditions, the study examined the filtration rate, growth and survival of four economically and ecologically important bivalve species native to the northeastern United States. They are the Eastern oyster (*Crassostrea virginica*); Atlantic surfclam (*Spisula solidissima*); hard clam (*Mercenaria mercenaria*); and softshell clam (*Mya arenaria*).

When supplied with seawater containing naturally occurring algal

particles, the groups that contained all four species removed significantly more particles than most monocultures. This suggests that each species prefers to filter a particular set of algal food particles.

"This shows that, to some degree, these bivalve species complement each other," said co-author Daphne Munroe, an associate professor in the Department of Marine and Coastal Sciences in the School of Environmental and Biological Sciences. She is based at the Haskin Shellfish Research Laboratory.

The scientists also found virtually no differences in growth or survival for any of the four species, suggesting that when food is not an issue, these bivalves could be raised together without outcompeting each another.

"This study illustrates the benefits of diversifying crops on shellfish farms," Acquafredda said. "Crop diversification gives aquaculture farmers protection from any individual crop failure, whether it's due to disease, predation or fluctuating environmental conditions. In future studies, the feasibility of bivalve polyculture should be tested on commercial bivalve farms."

**More information:** MP Acquafredda et al, The effect of species diversity on particle clearance and productivity in farmed bivalves, *Marine Ecology Progress Series* (2020). [DOI: 10.3354/meps13266](https://doi.org/10.3354/meps13266)

Provided by Rutgers University

Citation: Oysters and clams can be farmed together (2020, April 2) retrieved 24 May 2024 from <https://phys.org/news/2020-04-oysters-clams-farmed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.