

# **Wild tomato resistance to bacterial canker has implications for commercial tomato industry**

April 7 2020

---



F. Christopher Peritore-Galve, lead author of paper, and fellow graduate student Martha Sudermann. Credit: F. Christopher Peritore-Galve

Bacterial canker of tomato is a disease that leads to wilt, cankers, and eventually death. The disease was first discovered in Grand Rapids, Michigan, in 1909, but annual outbreaks now affect tomato production

areas worldwide. For some farmers, bacterial canker can be devastating and spoil an entire season's planting.

Bacterial canker is caused by the pathogen *Clavibacter michiganensis*, which infects commercially bred tomatoes by colonizing the xylem, a series of tubes that transports water and minerals throughout the plant. There are no commercially available tomatoes resistant to bacterial canker and management options are limited. However, breeders have known that wild tomato species are less susceptible to bacterial canker, but this knowledge is limited.

Scientists at the School of Integrative Plant Science at Cornell University were interested in understanding how *C. michiganensis* colonized the xylem in wild tomato and comparing to colonization of cultivated tomatoes. They found a [significant difference](#), according to the paper published in *Phytopathology*.

F. Christopher Peritore-Galve, lead author of this paper, explains: "Interestingly, the pathogen colonizes to high densities close to the inoculation site in wild tomatoes, but it is found in much lower densities 5 and 10 centimeters above that site, meaning that the bacteria are unable to spread through the xylem as well as in cultivated tomato xylem."

Additional research showed that the pathogen was restricted to the protoxylem vessels in vascular bundles of wild tomato.

"Protoxylem are early formed xylem vessels that are smaller in diameter that enable [water transport](#) during early, rapid plant growth," said Peritore-Galve. "We currently do not know why *C. michiganensis* preferentially colonizes these vessels, but that will be an avenue of future research."

Understanding what makes wild tomatoes able to tolerate this pathogen may help with future breeding efforts. For more information about this study, read "[Characterizing Colonization Patterns of \*Clavibacter michiganensis\* During Infection of Tolerant Wild \*Solanum\* Species.](#)"

**More information:** F. Christopher Peritore-Galve et al, Characterizing Colonization Patterns of *Clavibacter michiganensis* During Infection of Tolerant Wild *Solanum* Species, *Phytopathology*<sup>TM</sup> (2019). [DOI: 10.1094/PHYTO-09-19-0329-R](#)

Provided by American Phytopathological Society

Citation: Wild tomato resistance to bacterial canker has implications for commercial tomato industry (2020, April 7) retrieved 25 April 2024 from <https://phys.org/news/2020-04-wild-tomato-resistance-bacterial-canker.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.