


# Asteroids Ryugu and Bennu were formed by the destruction of a large asteroid

June 1 2020, by Brittany Enos

---



Sequence of images showing the formation of an aggregate by the reaccumulation of fragments produced during the disruption of an asteroid. Its final shape, five hours after the beginning of the process, is similar to that of Bennu and Ryugu. Credit: Michel et al./Nature Communications

Scientists with NASA's first asteroid sample return mission, OSIRIS-REx, are gaining a new understanding of asteroid Bennu's carbon-rich material and signature "spinning-top" shape. The team, led by the University of Arizona, has discovered that the asteroid's shape and hydration levels provide clues to the origins and histories of this and other small bodies.

Bennu, the target asteroid for the OSIRIS-REx mission, and Ryugu, the target of the Japan Aerospace Exploration Agency's Hayabusa2 asteroid

sample return mission, are composed of fragments of larger bodies that shattered upon colliding with other objects. The small fragments reaccumulated to form an aggregate body. Bennu and Ryugu may actually have formed in this way from the same original shattered parent body. Now, scientists are looking to discover what processes led to specific characteristics of these asteroids, such as their shape and mineralogy.

Bennu and Ryugu are both classified as "spinning-top" asteroids, which means they have a pronounced equatorial ridge. Until now, scientists thought that this shape formed as the result of thermal forces, called the YORP effect. The YORP effect increases the speed of the asteroid's spin, and over millions of years, material near the poles could have migrated to accumulate on the equator, eventually forming a spinning-top shape—meaning that the shape would have formed relatively recently.

However, in a new paper published in *Nature Communications*, scientists from the OSIRIS-REx and Hayabusa2 teams argue that the YORP effect may not explain the shape of either Bennu or Ryugu. Both asteroids have large impact craters on their equators, and their size suggests that these craters are some of Bennu's oldest surface features. Since the craters cover the equatorial ridges, their spinning-top shapes must also have been formed much earlier.

"Using computer simulations that model the impact that broke up Bennu's parent body, we show that these asteroids either formed directly as top-shapes, or achieved the shape early after their formation in the main [asteroid belt](#)," said Ronald Ballouz, co-lead author and OSIRIS-REx postdoctoral research associate at the UArizona. "The presence of the large equatorial craters on these asteroids, as seen in images returned by the spacecraft, rules out that the asteroids experienced a recent re-shaping due to the YORP effect. We would expect these craters to have

disappeared with a recent YORP-induced re-shaping of the asteroid."

In addition to their shapes, Bennu and Ryugu also both contain water-bearing surface material in the form of clay minerals. Ryugu's surface material is less water-rich than Bennu's, which implies that Ryugu's material experienced more heating at some point.

Assuming Bennu and Ryugu formed simultaneously, the paper explores two possible explanations for the different hydration levels of the two bodies based on the team's computer simulations. One hypothesis suggests that when the parent asteroid was disrupted, Bennu formed from material closer to the original surface, while Ryugu contained more material from near the parent body's original center. Another possible explanation for the difference in hydration levels is that the fragments experienced different levels of heating during the parent asteroid's [disruption](#). If this is the case, Ryugu's source material is likely from an area near the impact point, where temperatures were higher. Bennu's material would have come from a region that didn't undergo as much heating, and was likely farther from the point of impact. Analysis of the returned samples and further observational analysis of the asteroids' surfaces will provide a clearer idea of the possible shared history of the two asteroids.

"These simulations provide valuable new insights into how Bennu and Ryugu formed," said Dante Laretta, OSIRIS-REx principal investigator and UArizona professor of planetary sciences. "Once we have the returned samples of these two asteroids in the lab, we may be able to further confirm these models, possibly revealing the true relationship between the two asteroids."

Scientists anticipate that the samples will also provide new insights into the origins, formation and evolution of other carbonaceous asteroids and meteorites. The Japan Aerospace Exploration Agency's Hayabusa2

mission is currently making its way back to Earth, and is scheduled to deliver its samples of Ryugu late this year. The OSIRIS-REx mission will perform its first sample collection attempt at Bennu on Oct. 20 and will deliver its samples to Earth on Sep. 24, 2023.

**More information:** P. Michel et al, Collisional formation of top-shaped asteroids and implications for the origins of Ryugu and Bennu, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-16433-z](https://doi.org/10.1038/s41467-020-16433-z)

Provided by University of Arizona

Citation: Asteroids Ryugu and Bennu were formed by the destruction of a large asteroid (2020, June 1) retrieved 17 June 2024 from <https://phys.org/news/2020-05-asteroids-ryugu-bennu-destruction-large.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.