

# The carnivorous plant lifestyle is gene costly

May 15 2020

---



The genomes of the carnivorous plants Venus flytrap, spoon-leaved sundew and waterwheel (from left) are decoded. Credit: Dirk Becker and Sönke Scherzer / University of Würzburg

Plants can produce energy-rich biomass with the help of light, water and carbon dioxide. This is why they are at the beginning of the food chains. But carnivorous plants have turned the tables and prey on animals. Insects are their main food source.

A publication in the journal *Current Biology* now sheds light on the secret life of the green carnivores. The plant scientist Rainer Hedrich and the evolutionary bioinformatician Jörg Schultz, both from Julius-Maximilians-Universität (JMU) Würzburg in Bavaria, Germany, and

their colleague Mitsujasu Hasebe from the University of Okazaki (Japan) have deciphered and analysed the genomes of three carnivorous plant species.

They studied the Venus flytrap *Dionaea muscipula*, which originates from North America, the globally occurring waterwheel plant *Aldrovanda vesiculosa*, and the spoon-leaved sundew *Drosera spatulata*, which is widely distributed in Asia.

All three belong to the sundew family. Nevertheless, they have each conquered different habitats and developed their own trapping mechanisms. In *Dionaea* and *Aldrovanda*, the ends of the leaves are transformed into folding traps. The sundew, on the other hand, attaches its prey to the leaf surface with sticky tentacles.

## **Basic genes for carnivory**

The first thing the international research team found out was that, despite their different lifestyles and trapping mechanisms, Venus flytrap, sundew and waterwheel have a common "basic set" of genes that are essential for the carnivorous lifestyle.

"The function of these genes is related to the ability to sense and digest prey animals and to utilise their nutrients," explains Rainer Hedrich.

"We were able to trace the origin of the carnivory genes back to a duplication event that occurred many millions of years ago in the genome of the last common ancestor of the three carnivorous species," says Jörg Schultz. The duplication of the entire genome has provided evolution with an ideal playing ground for developing new functions.

## **Genetic poverty despite a special way of life**

To their surprise, the researchers discovered that the [plants](#) do not need a particularly large number of genes for carnivory. Instead, the three species studied are actually among the most gene-poor plants known. Drosera has 18,111, Dionaea 21,135 and Aldrovanda 25,123 genes. In contrast, most plants have between 30,000 and 40,000 genes.

How can this be reconciled with the fact that a wealth of new genes is usually needed to develop new ways of life? "This can only mean that the specialization in animal food was accompanied by an increase in the number of genes, but also a massive loss of genes," concludes developmental biologist Hasebe.

## **Root genes are active in the trapping organs**

Most of the genes required for the insect traps are also found in slightly modified form in normal plants. "In [carnivorous plants](#), several genes are active in the trapping organs, which in other plants have their effect in the root. In the trapping organs, these [genes](#) are only switched on when the prey is secure," explains Hedrich. This finding is consistent with the fact that the roots are considerably reduced in Venus flytrap and sundew. In the waterwheel they are completely absent.

## **Further research into the trapping function**

The researchers now have an insight into the evolution of carnivory in plants and hold three blueprints for this particular way of life in their hands. Their next goal is to gain an even better understanding of the molecular basis of the trapping function.

"We have found that the Venus flytrap counts the electrical stimuli triggered by the prey, can remember this number for a certain time and finally makes a decision that corresponds to the number," says Hedrich.

Now it is important to understand the biophysical-biochemical principle according to which carnivorous plants count.

**More information:** Gergo Palfalvi et al, Genomes of the Venus Flytrap and Close Relatives Unveil the Roots of Plant Carnivory, *Current Biology* (2020). [DOI: 10.1016/j.cub.2020.04.051](https://doi.org/10.1016/j.cub.2020.04.051)

Provided by University of Würzburg

Citation: The carnivorous plant lifestyle is gene costly (2020, May 15) retrieved 25 April 2024 from <https://phys.org/news/2020-05-carnivorous-lifestyle-gene-costly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.