

Chirality-assisted lateral momentum transfer for bidirectional enantioselective separation

May 26 2020





a Illustration of the sorting of chiral particles with left and right handedness by the optical lateral force at an air-water interface. b Microscopic image of the chiral microparticles. The scale bar equals 5 μ m. c SEM image of dry polymeric chiral microparticles. The scale bar equals 1 μ m. d TEM images of thin slices of microparticles with different sizes and different spiral configurations. The scale bars equal 1 μ m. Electric field and 2D Poynting vector in the y-z plane for θ of 10° (e) and 45° (f). The Poynting vector strings has a bias to the left (e) and right (f), resulting positive (e) and negative (f) optical lateral forces respectively Credit: by Cheng-Wei Qiu

Light carrying photon momentum can push and pull microparticles through momentum exchange. This momentum exchange process generates optical forces, which either attracts (conventional optical tweezers), pushes (radiation force) or pulls (pulling force) microparticles. A new emerging research interest, optical lateral force which represents the optical force perpendicular to the propagating direction of a non-gradient beam, has attracted much attention. The lateral force can be generated using achiral particles through the conversion of spin and orbital momentum of a circularly polarized beam. It is also predicted that a chiral nanoparticle placed above a surface can generate the lateral force using a plane wave excitation. However, there are few demonstrations of the chirality dependent lateral force, and the particles used in the theoretical prediction are 100 nm which has limited applications. Besides, the theory of optical lateral force on bigger particles (size ~ wavelength) is lacking.

In a new paper published in *Light Science & Applications*, scientists designed an experiment to demonstrate the chirality-dependent optical <u>lateral force</u>. They synthesize microparticles with strong chirality and float them at the interface of air and water. After illumination with an obliquely incident linearly polarized beam, chiral microparticles with



different handedness (left and right) will move in opposite directions. Interestingly, they find in theory that the optical lateral <u>force</u> could reverse signs with different light polarization and incident angle, chirality value and particle size. They also developed intriguing models in the perspective of momentum transfer to elaborate this optical lateral force. The reported method and technique will open new avenues for future direct detection and sorting of microparticles with imperceptible chemical differences and inspire the exploration of optical phenomena with light-matter interactions.

The experimental setup is facile, only requiring an s- or p-polarized laser beam and focusing it into an elliptical shape using two cylindrical lenses. The chiral microparticles are floating at the interface of air and water in a microwell fabricated using soft lithography. This configuration offers many possibilities to probe intriguing optical physics, such as spinorbital interactions, chirality sensing, etc.





a, b Chiral particles with right hand ($\kappa > 0$) (a) and left hand (κ

These scientists summarize their work as:

"We designed the first experiment of chirality assisted optical lateral force on Mie chiral particles (size ~ wavelength) for enantioselective separation. Recent studies about enantioselective separation and chiral particles focus on the very big particles (geometric optics region, size > wavelength) due to the limit of particle synthesis procedure and existing theories. We demonstrated, for the first time, robust bidirectional sorting of Mie chiral particles, and the first example of



reversible optical lateral forces, which we believe is an essential complement to the community of optics as well as optical manipulations. Our theory studies the dependence of the reversible optical lateral forces with <u>particle size</u>, incident angle, and polarization of light. Compared to previous non-reversal optical lateral force, Mie chiral particles are quite unique and nontrivial, and they have some intriguing properties. Also, we elaborate on the optical lateral force from the perspective of momentum transfer, which is a straightforward way to manifest the optical lateral force."

"Our method is insightful and helpful to the demonstration of extraordinary forces since it rules out the optical gradient forces in conventional optical tweezers. It helps to complement the realm of optical lateral forces in both theory and experiment. The presented technique can be used for contactless monitoring of the handedness of the chiral particles which exist widely in the drug industry and biomaterials without testing with chemical or biological methods," Dr. Yuzhi Shi and Prof. Cheng-Wei Qiu added.

More information: Yuzhi Shi et al, Chirality-assisted lateral momentum transfer for bidirectional enantioselective separation, *Light: Science & Applications* (2020). DOI: 10.1038/s41377-020-0293-0

Provided by Chinese Academy of Sciences

Citation: Chirality-assisted lateral momentum transfer for bidirectional enantioselective separation (2020, May 26) retrieved 1 May 2024 from <u>https://phys.org/news/2020-05-chirality-assisted-lateral-momentum-bidirectional-enantioselective.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.