

# Dinosaur-dooming asteroid struck earth at 'deadliest possible' angle

May 26 2020, by Caroline Brogan

---



Original artwork depicting the moment the asteroid struck in present-day Mexico. Credit: Chase Stone

New simulations from Imperial College London have revealed the

asteroid that doomed the dinosaurs struck Earth at the 'deadliest possible' angle.

The simulations show that the asteroid hit Earth at an angle of about 60 degrees, which maximised the amount of climate-changing gases thrust into the [upper atmosphere](#).

Such a strike likely unleashed billions of tonnes of sulphur, blocking the sun and triggering the nuclear winter that killed the dinosaurs and 75 per cent of life on Earth 66 million years ago.

Drawn from a combination of 3-D numerical impact simulations and [geophysical data](#) from the site of the impact, the new models are the first ever fully 3-D simulations to reproduce the whole event—from the initial impact to the moment the final crater, now known as Chicxulub, was formed.

The simulations were performed on the Science and Technology Facilities Council (STFC) DiRAC High Performance Computing Facility.

Lead researcher Professor Gareth Collins, of Imperial's Department of Earth Science and Engineering, said: "For the dinosaurs, the [worst-case scenario](#) is exactly what happened. The asteroid strike unleashed an incredible amount of climate-changing gases into the atmosphere, triggering a chain of events that led to the extinction of the dinosaurs. This was likely worsened by the fact that it struck at one of the deadliest possible angles.

"Our simulations provide compelling evidence that the asteroid struck at a steep angle, perhaps 60 degrees above the horizon, and approached its target from the north-east. We know that this was among the worst-case scenarios for the lethality on impact, because it put more hazardous

debris into the upper atmosphere and scattered it everywhere—the very thing that led to a nuclear winter."

The results are published today in *Nature Communications*.

## **Crater creation**

The upper layers of earth around the Chicxulub crater in present-day Mexico contain high amounts of water as well as porous carbonate and evaporite rocks. When heated and disturbed by the impact, these rocks would have decomposed, flinging vast amounts of carbon dioxide, sulphur and water vapour into the atmosphere.

The sulphur would have been particularly hazardous as it rapidly forms aerosols—tiny particles that would have blocked the sun's rays, halting photosynthesis in plants and rapidly cooling the climate. This eventually contributed to the mass extinction event that killed 75 per cent of life on Earth.

The team of researchers from Imperial, the University of Freiburg, and The University of Texas at Austin, examined the shape and subsurface structure of the crater using geophysical data to feed into the simulations that helped diagnose the impact angle and direction. Their analysis was also informed by recent results from drilling into the 200 km-wide crater, which brought up rocks containing evidence of the extreme forces generated by the impact.

## **Peak performance**

Pivotal to diagnosing the angle and direction of impact was the relationship between the centre of the crater, the centre of the peak ring—a ring of mountains made of heavily fractured rock inside the

crater rim—and the centre of dense uplifted mantle rocks, some 30 km beneath the crater.

At Chicxulub, these centres are aligned in a southwest-northeast direction, with the crater centre in between the peak-ring and mantle-uplift centres. The team's 3-D Chicxulub crater simulations at an angle of 60 degrees reproduced these observations almost exactly.

The simulations reconstructed the crater formation in unprecedented detail and give us more clues as to how the largest craters on Earth are formed. Previous fully 3-D simulations of the Chicxulub impact have covered only the early stages of impact, which include the production of a deep bowl-shaped hole in the crust known as the transient crater and the expulsion of rocks, water and sediment into the atmosphere.

These simulations are the first to continue beyond this intermediate point in the formation of the crater and reproduce the final stage of the crater's formation, in which the transient crater collapses to form the final structure (see video). This allowed the researchers to make the first comparison between 3-D Chicxulub crater simulations and the present-day structure of the crater revealed by geophysical data.

Co-author Dr. Auriol Rae of the University of Freiburg said: "Despite being buried beneath nearly a kilometre of sedimentary rocks, it is remarkable that geophysical data reveals so much about the crater structure—enough to describe the direction and angle of the impact."

The researchers say that while the study has given us important insights into the dinosaur-dooming impact, it also helps us understand how large craters on other planets form.

Co-author Dr. Thomas Davison, also of Imperial's Department of Earth Science and Engineering, said: "Large craters like Chicxulub are formed

in a matter of minutes, and involve a spectacular rebound of rock beneath the [crater](#). Our findings could help advance our understanding of how this rebound can be used to diagnose details of the impacting asteroid."

**More information:** Collins, G.S., Patel, N., Davison, T.M. et al. A steeply-inclined trajectory for the Chicxulub impact. *Nat Commun* 11, 1480 (2020). [doi.org/10.1038/s41467-020-15269-x](https://doi.org/10.1038/s41467-020-15269-x)

Provided by Imperial College London

Citation: Dinosaur-dooming asteroid struck earth at 'deadliest possible' angle (2020, May 26) retrieved 20 March 2024 from <https://phys.org/news/2020-05-dinosaur-dooming-asteroid-struck-earth-deadliest.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--