

Newly modeled: Minimum energy requirements for microbial communities to live

May 6 2020, by Alice Scott



Credit: CC0 Public Domain

A microbial community is a complex, dynamic system composed of



hundreds of species and their interactions, they are found in oceans, soil, animal guts and plant roots. Each system feeds the Earth's ecosystem and their own growth, as they each have their own metabolism that underpin biogeochemical cycles.

The same community-level <u>metabolic rates</u> are exploited in biotechnology for water treatment and bioenergy production from organic waste, thus the ability to capture microbial growth rates and metabolic activities within the communities is key for modeling of planetary ecosystem dynamics, animal and plant health and biotechnological waste valorzation.

Models of such systems should account for both kinetic and thermodynamic constraints inherent in microbial growth, which is a challenge due to the complexity of these systems.

However in the paper, "Thermodynamic modeling of synthetic communities predicts minimum free energy requirements for sulfate reduction and methanogenesis," published 6 May in the *Journal of the Royal Society Interface*, researchers from the School of Life Sciences at the University of Warwick have produced an extendable thermodynamic model for simulating the dynamics of microbial communities.

To develop and calibrate a thermodynamic model of microbial growth and metabolite dynamics in a microbial context, researchers focused on defined anaerobic synthetic communities—these communities breakdown biodegradable material in the absence of oxygen.

They used a recently developed experimental model system for studying syntrophic interactions among sulfate reducers and methanogens, which make up a key part of anaerobic <u>microbial communities</u> found in reactors and freshwater and estuary sediments.



Researchers found the minimum energy requirement for a given <u>metabolic pathway</u>, providing evidence that experimental data can be used to estimate energy requirements of microbial pathways, and that such estimates for methanogenesis (the production of methane) and sulfate reduction.

Professor Orkun Soyer, from the School of Life Sciences at the University of Warwick comments:

"Microbes are crucial in mediating biochemical conversions in the environment. These conversions ultimately allow higher organisms like us and plants to obtain the metabolites that they need for living, and underpin biogeochemical cycles that make Earth a habitable planet. Better understanding these processes requires modeling the biochemical conversions mediated by microbes.

"Our recent work creates a generalizable platform for this purpose and introduces a thermodynamic model of microbial conversions. This <u>model</u> better captures dynamics of metabolic conversions and provides estimates of minimal energy requirements of such conversions from <u>experimental data</u>."

More information: Hadrien Delattre et al. Thermodynamic modelling of synthetic communities predicts minimum free energy requirements for sulfate reduction and methanogenesis, *Journal of The Royal Society Interface* (2020). DOI: 10.1098/rsif.2020.0053

Provided by University of Warwick

Citation: Newly modeled: Minimum energy requirements for microbial communities to live (2020, May 6) retrieved 3 May 2024 from <u>https://phys.org/news/2020-05-newly-minimum-</u>



energy-requirements-microbial.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.