


Next-gen laser facilities look to usher in new era of relativistic plasmas research

May 26 2020



Quantum electrodynamics phenomena in plasmas. Credit: Stephen Alvey/Alec Thomas

The subject of the 2018 Nobel Prize in physics, chirped pulse amplification is a technique that increases the strength of laser pulses in many of today's highest-powered research lasers. As next-generation laser facilities look to push beam power up to 10 petawatts, physicists expect a new era for studying plasmas, whose behavior is affected by features typically seen in black holes and the winds from pulsars.

Researchers released a study taking stock of what upcoming high-power

laser capabilities are poised to teach us about relativistic plasmas subjected to strong-field quantum electrodynamics (QED) processes. In addition, the proposed new study designs for further exploring these new phenomena.

Appearing in *Physics of Plasmas*, the article introduces the physics of relativistic [plasma](#) in supercritical fields, discusses the current state of the field and provides an overview of recent developments. It also highlights open questions and topics that are likely to dominate the attention of people working in the field over the next several years.

Strong-field QED is a lesser-studied corner of the standard model of particle physics that has not been explored at big collider facilities, such as SLAC National Accelerator Laboratory or CERN, the European Organization for Nuclear Research, due to the lack of strong electromagnetic fields in accelerator settings. With high-intensity lasers, researchers can use strong fields, which have been observed in phenomena such as gamma ray emission and electron-positron pair production.

The group explores how the findings could potentially lead to advances in studies of fundamental physics and in the development of high-energy ion, electron, positron and photon sources. Such findings would be crucial for expanding on many types of scanning technology present today, ranging from materials science studies to medical radiotherapy to next-generation radiography for homeland security and industry.

The QED processes will result in dramatically new plasma physics phenomena, such as the generation of dense electron-positron pair plasma from near vacuum, complete laser energy absorption by QED processes, or the stopping of an ultrarelativistic electron beam, which could penetrate a centimeter of lead by a hair's breadth of [laser](#) light.

"What kind of new technology these new plasma physics phenomena might translate is largely unknown, especially because the field of QED plasmas itself is a kind of uncharted territory in [physics](#)," author Peng Zhang said. "At the current stage, even adequate theoretical understanding is significantly lacking."

The group hopes the paper will help bring the attention of more researchers to the exciting new fields of QED plasmas.

More information: "Relativistic plasma physics in supercritical fields," *Physics of Plasmas* (2020).

aip.scitation.org/doi/10.1063/1.5144449

Provided by American Institute of Physics

Citation: Next-gen laser facilities look to usher in new era of relativistic plasmas research (2020, May 26) retrieved 3 May 2024 from

<https://phys.org/news/2020-05-next-gen-laser-facilities-usher-era.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.