

# Permafrost in the Arctic can thaw faster than presumed

May 26 2020

---



A thermokarst pond in Siberia. Massive ground ice layers are visible under the vegetation. When they melt, the surface can collapse into depressions. According to the new study, such processes could dominate the landscape in the future.

Credit: Jan Nitzbon

Melting of ice in permafrost ground leads to processes of change in the landscape—thermokarst. This may cause faster thawing of the permafrost.

Some of the coldest [permafrost](#) on Earth might be more vulnerable to thawing than previously thought. In the PERMANOR project led by the University of Oslo in Norway, an international research team has investigated so-called thermokarst processes which can significantly accelerate thawing of permafrost. The study is recently published in *Nature Communications*.

### **Reinforces warming**

Air temperatures are increasing in [high latitudes](#) and in high mountain areas dominated by permafrost in the ground. When this [permafrost thaws](#), large amounts of greenhouse gases can be released to the atmosphere, which can potentially intensify [global warming](#).

Another consequence is that ice layers in the ground start to melt, so that the ground subsides and depressions with ponds and lakes form. This [landscape](#) change, which is known as "thermokarst", can once again accelerate permafrost thawing in a positive feedback to global [warming](#).

### **Changes the landscape**

The research team from Germany and Norway has developed computer models to investigate how thermokarst processes affect the future of permafrost in Northeastern Siberia. Today this region is dominated by very cold and supposedly stable permafrost.

So far, thermokarst processes have not been included in models that simulate the permafrost evolution in a warming climate. "We found that when including thermokarst processes in our simulations, even cold

permafrost in Northern Siberia could thaw at the end of this century, although the ground currently still has temperatures of minus ten degrees" says lead author Jan Nitzbon from the Alfred Wegener Institute for Polar and Marine Research in Germany.



Thawing permafrost and thermokarst processes have made ponds in this peatland landscape in Northern Norway. In the PERMANOR project these processes are systematically mapped with a drone over several years. The small pond in the foreground is only a few months old, while the other has grown considerably during the last four years. Credit: Sebastian Westermann

Among scientists, the general assumption is that thermokarst processes locally lead to faster thawing of permafrost.

"However, with our model, we have also considered stabilizing processes

that can slow down thawing. We were amazed that under a moderate warming scenario, thermokarst processes can even limit the thawing of permafrost", he states to Titan.uio.no.

Under a stronger warming scenario, however, self-reinforcing processes dominated, which would drastically change these landscapes due to accelerated permafrost thaw.

### **Also applies to permafrost in Northern Norway**

Sebastian Westermann, Associate Professor from the University of Oslo who has led the PERMANOR project, adds:

"Thermokarst not only affects permafrost regions in Siberia, we see the same processes in Norway, especially in peatlands in Finnmark. Within just a few years, new ponds have formed in our study areas, decreasing the permafrost area. In Norway, the permafrost is already now close to the freezing point, and we are observing many of the landscape changes that could occur in Siberia in the future."

### **Improves global climate model**

The researchers have now started to implement the new simulation tools in Earth System Models, such as the Norwegian climate [model](#) NorESM. With this, it will be possible to improve future simulations of permafrost thaw not only in Siberia, but on the global scale.

The results of the present study strongly suggest that limiting global warming can help protect arctic ecosystems and landscapes from drastic changes due to thawing of ice-rich permafrost.

**More information:** Jan Nitzbon et al. Fast response of cold ice-rich permafrost in northeast Siberia to a warming climate, *Nature*

*Communications* (2020). [DOI: 10.1038/s41467-020-15725-8](https://doi.org/10.1038/s41467-020-15725-8)

Provided by University of Oslo

Citation: Permafrost in the Arctic can thaw faster than presumed (2020, May 26) retrieved 24 March 2023 from <https://phys.org/news/2020-05-permafrost-arctic-faster-presumed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.