


Mapping dry wildfire fuels with AI and new satellite data

May 21 2020, by Josie Garthwaite



Examples of forest dryness progressing across western states in 2019. Credit: Krishna Rao

As California and the American West head into fire season amid the coronavirus pandemic, scientists are harnessing artificial intelligence and new satellite data to help predict blazes across the region.

Anticipating where a fire is likely to ignite and how it might spread requires information about how much burnable plant material exists on the landscape and its dryness. Yet this information is surprisingly difficult to gather at the scale and speed necessary to aid wildfire

management.

Now, a team of experts in hydrology, remote sensing and [environmental engineering](#) have developed a deep-learning model that maps [fuel](#) moisture levels in fine detail across 12 western states, from Colorado, Montana, Texas and Wyoming to the Pacific Coast.

The researchers describe their technique in the August 2020 issue of Remote Sensing of Environment. According to the senior author of the paper, Stanford University ecohydrologist Alexandra Konings, the new dataset produced by the model could "massively improve fire studies."

According to the paper's lead author, Krishna Rao, a Ph.D. student in Earth system science at Stanford, the model needs more testing to figure into fire management decisions that put lives and homes on the line. But it's already illuminating previously invisible patterns. Just being able to see forest dryness unfold pixel by pixel over time, he said, can help reveal areas at greatest risk and "chart out candidate locations for prescribed burns."

The work comes at a time of growing urgency for this kind of insight, as climate change extends and intensifies the wildfire season—and as the ongoing COVID-19 pandemic complicates efforts to prevent large fires through controlled burns, prepare for mass evacuations and mobilize first responders.




Smoke from the 2016 Cedar Fire rises above trees in Sequoia National Forest.
Credit: Lance Cheung/USDA

Getting a read on parched landscapes

Fire agencies today typically gauge the amount of dried-out, flammable vegetation in an area based on samples from a small number of trees. Researchers chop and weigh tree branches, dry them out in an oven and then weigh them again. "You look at how much mass was lost in the oven, and that's all the water that was in there," said Konings, an assistant professor of Earth system science in Stanford's School of Earth, Energy & Environmental Sciences (Stanford Earth). "That's obviously really laborious, and you can only do that in a couple of different places, for only some of the species in a landscape."

The U.S. Forest Service painstakingly collects this plant water content data at hundreds of sites nationwide and adds them to the National Fuel Moisture Database, which has amassed some 200,000 such measurements since the 1970s. Known as live fuel moisture content, the metric is well established as a factor that influences wildfire risk. Yet little is known about how it varies over time from one plant to another—or from one ecosystem to another.

For decades, scientists have estimated fuel moisture content indirectly, from informed but unproven guesses about relationships between temperature, precipitation, water in dead plants and the dryness of living ones. According to Rao, "Now, we are in a position where we can go back and test what we've been assuming for so long—the link between weather and live fuel moisture—in different ecosystems of the western United States."



Maps display the amount of water in plants relative to dry biomass across the American West. Credit: Krishna Rao

AI with a human assist

The new model uses what's called a recurrent neural network, an artificial intelligence system that can learn to recognize patterns in vast

mountains of data. The scientists trained their model using field data from the National Fuel Moisture Database, then put it to work estimating fuel moisture from two types of measurements collected by spaceborne sensors. One involves measurements of visible light bouncing off Earth. The other, known as synthetic aperture radar (SAR), measures the return of microwave radar signals, which can penetrate through leafy branches all the way to the ground surface.

"One of our big breakthroughs was to look at a newer set of satellites that are using much longer wavelengths, which allows the observations to be sensitive to water much deeper into the forest canopy and be directly representative of the fuel moisture content," said Konings, who is also a center fellow, by courtesy, at Stanford Woods Institute for the Environment.

To train and validate the model, the researchers fed it three years of data for 239 sites across the American west starting in 2015, when SAR data from the European Space Agency's Sentinel-1 satellites became available. They checked its fuel moisture predictions in six common types of land cover, including broadleaf deciduous forests, needleleaf evergreen forests, shrublands, grasslands and sparse vegetation, and found they were most accurate—meaning the AI predictions most closely matched field measurements in the National Fuel Moisture Database—in shrublands.

Rich with aromatic herbs like rosemary and oregano, and often marked by short trees and steep, rocky slopes, shrublands occupy as much as 45 percent of the American West. They're not only the region's biggest ecosystem, Rao said, "they are also extremely susceptible to frequent fires since they grow back rapidly." In California, fires whipped to enormous size by Santa Ana winds burn in a type of shrubland known as chaparral. "This has led fire agencies to monitor them intensively," he said.

The model's estimates feed into an [interactive map](#) that fire agencies may eventually be able to use to identify patterns and prioritize control measures. For now, the map offers a dive through history, showing fuel moisture content from 2016 to 2019, but the same method could be used to display current estimates. "Creating these maps was the first step in understanding how this new fuel [moisture](#) data might affect fire risk and predictions," Konings said. "Now we're trying to really pin down the best ways to use it for improved [fire](#) prediction."

More information: Krishna Rao et al. SAR-enhanced mapping of live fuel moisture content, *Remote Sensing of Environment* (2020). [DOI: 10.1016/j.rse.2020.111797](#)

Provided by Stanford University

Citation: Mapping dry wildfire fuels with AI and new satellite data (2020, May 21) retrieved 24 April 2024 from <https://phys.org/news/2020-05-wildfire-fuels-ai-satellite.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--