

ATLAS experiment finds evidence of spectacular four-top quark production

June 23 2020, by Atlas Collaboration,



Figure 1: Event display of a candidate four-top-quark event, where two of the top quarks decay leptonically (one with a resulting muon (red) and one with an electron (green)), and two top quarks decay hadronically (green and yellow rectangles). The jets (b-tagged jets) are shown as yellow (blue) cones. Credit: ATLAS Collaboration/CERN



The ATLAS Collaboration at CERN has announced strong evidence of the production of four top quarks. This rare Standard Model process is expected to occur only once for every 70 thousand pairs of top quarks created at the Large Hadron Collider (LHC) and has proven extremely difficult to measure.

The top quark is the most massive elementary particle in the Standard Model, clocking in at 173 GeV, which is equivalent to the mass of a gold atom. But contrary to gold, whose mass is mostly due to the nuclear binding force, the top quark gets all its mass from the interaction with the Higgs field. So when four top quarks are produced in a single event, they create the heaviest particle final state ever seen at the LHC, with almost 700 GeV in total. This is an ideal environment to search for new physics with yet unknown particles contributing to the process. Should they exist, physicists will see additional production of four top quarks above what is predicted by the Standard Model, further motivating a detailed study of the process.

In their new search for four-top-quark production, ATLAS physicists studied the full Run 2 dataset recorded between 2015 and 2018. When produced through proton–proton collisions at the LHC, this process leaves spectacular signatures in the ATLAS detector. The four top quarks produce four W bosons and four jets—collimated sprays of particles—originating from bottom quarks. The W bosons then, in turn, each decay into two jets or one charged lepton (electron, muon or tau leptons) and an invisible neutrino. As a final step, the tau leptons decay into a lighter lepton or a jet, with additional neutrinos.

For this result, physicists chose to focus on collision events producing two leptons with the same charge or three leptons. Despite accounting for only 12% of all four-top-quark decays, these signatures are easier to distinguish from background processes in the ATLAS detector. Detecting a signal nevertheless required a detailed understanding of the



remaining background processes and the use of sophisticated separation techniques.



Figure 2: The boosted decision tree (BDT) score output for the signal region (SR). The data are shown in black; the simulated signal in red. The y-axis shows



the number of events and is in the logarithmic-scale. The band includes the total uncertainty on the post profile-likelihood fit (post-fit) computation. The ratio of the data to the total post-fit computation is shown in the lower panel. Credit: ATLAS Collaboration/CERN

ATLAS physicists trained a multivariate discriminant (boosted decision tree) using the distinct features of the signal, including the high numbers of jets, their quark-flavor origin (bottom quark or not), and the energies and angular distributions of the measured particles. The main background processes that resemble the signal stem from the production of a pair of top quarks in association with other particles, such as a W or Z boson, a Higgs boson, or another top quark. Some of these processes have themselves only recently been observed by the ATLAS and CMS Collaborations.

Each background process was individually evaluated, primarily through dedicated simulations which included information from the best available theoretical predictions. The most difficult background processes—the top-quark-pair production with a W boson and backgrounds with fake leptons—had to be determined using data from dedicated control regions. Fake leptons arise when the charge of a lepton is misidentified, or when leptons come from a different process, but are attributed to the signal. Both had to be well-understood and precisely evaluated in order to reduce the systematic uncertainty on the final result.

ATLAS measured the cross section for the production of four top quarks to be 24 $^{+7}_{-6}$ fb, which is consistent with the Standard Model prediction (12 fb) at 1.7 standard deviations. The signal significance amounts to 4.3 standard deviations, for an expected significance of 2.4 standard deviations were the four-<u>top-quark</u> signal equal to the Standard Model



prediction. The measurement provides strong evidence for this process.

Additional data from the next LHC run—along with further developments of the analysis techniques employed—will improve the precision of this challenging measurement.

More information: Evidence for tttt production in the multilepton final state in proton-proton collisions at 13 TeV with the ATLAS detector (ATLAS-CONF-2020-013): <u>atlas.web.cern.ch/Atlas/GROUPS</u>....<u>ATLAS-CONF-2020-013/</u>

Provided by ATLAS Experiment

Citation: ATLAS experiment finds evidence of spectacular four-top quark production (2020, June 23) retrieved 5 May 2024 from https://phys.org/news/2020-06-atlas-evidence-spectacular-four-top-quark.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.