


# **Excitons form superfluid in certain 2-D combos**

June 15 2020, by Mike Williams

---



Rice University theorists determined that certain combinations of weakly bound 2D materials let holes and electrons combine into excitons at the materials' ground state. That combination can lead them to condense into a superfluidlike phase. The discovery shows promise for electronic, spintronic and quantum computing applications. Credit: Yakobson Research Group/Rice University

Mixing and matching computational models of 2-D materials led scientists at Rice University to the realization that excitons—quasiparticles that exist when electrons and holes briefly bind—can be manipulated in new and useful ways.

The researchers identified a small set of 2-D compounds with similar atomic lattice dimensions that, when placed together, would allow excitons to form spontaneously. Generally, excitons happen when energy from light or electricity boosts electrons and holes into a higher state.

But in a few of the combinations predicted by Rice materials theorist Boris Yakobson and his team, excitons were observed stabilizing at the materials' ground state. According to their determination, these excitons at their lowest energy state could condense into a superfluidlike phase. The discovery shows promise for electronic, spintronic and quantum computing applications.

"The very word '[exciton](#)' means that electrons and holes 'jump up' into a higher energy," Yakobson said. "All cold systems sit in their lowest-possible energy states, so no excitons are present. But we found a realization of what seems a paradox as conceived by Nevill Mott 60 years ago: a material system where excitons can form and exist in the ground state."

The open-access study by Yakobson, graduate student Sunny Gupta and research scientist Alex Kutana, all of Rice's Brown School of Engineering, appears in *Nature Communications*.

After evaluating many thousands of possibilities, the team precisely modeled 23 bilayer heterostructures, their layers loosely held in alignment by weak van der Waals forces, and calculated how their band gaps aligned when placed next to each other. (Band gaps define the distance an electron has to leap to give a material its semiconducting

properties. Perfect conductors—metals or semimetals like graphene—have no [band gap](#).)

Ultimately, they produced phase diagrams for each combination, maps that allowed them to view which had the best potential for experimental study.

"The best combinations are distinguished by a lattice parameter match and, most importantly, by the special positions of the electronic bands that form a broken gap, also called type III," Yakobson said.

Conveniently, the most robust combinations may be adjusted by applying stress through tension, curvature or an external electric field, the researchers wrote. That could allow the phase state of the excitons to be tuned to take on the "perfect fluid" properties of a Bose-Einstein condensate or a superconducting BCS condensate.

"In a quantum condensate, bosonic particles at low temperatures occupy a collective quantum ground state," Gupta said. "That supports macroscopic quantum phenomena as remarkable as superfluidity and superconductivity."

"Condensate states are intriguing because they possess bizarre quantum properties and exist on an everyday scale, accessible without a microscope, and only low temperature is required," Kutana added.

"Because they are at the lowest possible energy state and because of their quantum nature, condensates cannot lose energy and behave as a perfect frictionless fluid.

"Researchers have been looking to realize them in various solid and gas systems," he said. "Such systems are very rare, so having two-dimensional materials among them would greatly expand our window into the quantum world and create opportunities for use in new, amazing

devices."

The best combinations were assemblies of heterostructure bilayers of antimony-tellurium-selenium with bismuth-tellurium-chlorine; hafnium-nitrogen-iodine with zirconium-nitrogen-chlorine; and lithium-aluminum-tellurium with bismuth-tellurium-iodine.

"Except for having similar lattice parameters within each pair, the chemistry compositions appear rather nonintuitive," Yakobson said. "We saw no way to anticipate the desired behavior without the painstaking quantitative analysis.

"One can never deny a chance to find serendipity—as Robert Curl said, chemistry is all about getting lucky—but sifting through hundreds of thousands of material combinations is unrealistic in any lab. Theoretically, however, it can be done."

**More information:** Sunny Gupta et al, Heterobilayers of 2-D materials as a platform for excitonic superfluidity, *Nature Communications* (2020). DOI: [10.1038/s41467-020-16737-0](https://doi.org/10.1038/s41467-020-16737-0)

Provided by Rice University

Citation: Excitons form superfluid in certain 2-D combos (2020, June 15) retrieved 23 May 2024 from <https://phys.org/news/2020-06-excitons-superfluid-d-combos.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--