

Hot stars are plagued by giant magnetic spots

June 1 2020



Astronomers using ESO telescopes have discovered giant spots on the surface of extremely hot stars hidden in stellar clusters, called extreme horizontal branch stars. This image shows an artist's impression of what one of these stars, and its giant whitish spot, might look like. The spot is bright, takes up a quarter of the star's surface and is caused by magnetic fields. As the star rotates, the spot on its surface comes and goes, causing visible changes in brightness. Credit: ESO/L. Calçada, INAF-Padua/S. Zaggia

Astronomers using European Southern Observatory (ESO) telescopes

have discovered giant spots on the surface of extremely hot stars hidden in stellar clusters. Not only are these stars plagued by magnetic spots, some also experience superflare events, explosions of energy several million times more energetic than similar eruptions on the Sun. The findings, published today in *Nature Astronomy*, help astronomers better understand these puzzling stars and open doors to resolving other elusive mysteries of stellar astronomy.

The team, led by Yazan Momany from the INAF Astronomical Observatory of Padua in Italy, looked at a particular type of star known as extreme horizontal branch [stars](#)—objects with about half the mass of the Sun but four to five times hotter. "These hot and small stars are special because we know they will bypass one of the final phases in the life of a typical star and will die prematurely," says Momany, who was previously a staff astronomer at ESO's Paranal Observatory in Chile. "In our Galaxy, these peculiar hot objects are generally associated with the presence of a close companion star."

Surprisingly, however, the vast majority of these extreme horizontal branch stars, when observed in tightly packed stellar groups called globular clusters, do not appear to have companions. The team's long-term monitoring of these stars, made with ESO telescopes, also revealed that there was something more to these mysterious objects. When looking at three different globular clusters, Momany and his colleagues found that many of the extreme horizontal branch stars within them showed regular changes in their brightness over the course of just a few days to several weeks.

"After eliminating all other scenarios, there was only one remaining possibility to explain their observed brightness variations," concludes Simone Zaggia, a study co-author from the INAF Astronomical Observatory of Padua in Italy and a former ESO Fellow: "these stars must be plagued by spots!"

Spots on extreme horizontal branch stars appear to be quite different from the dark sunspots on our own Sun, but both are caused by magnetic fields. The spots on these hot, extreme stars are brighter and hotter than the surrounding stellar surface, unlike on the Sun where we see spots as dark stains on the solar surface that are cooler than their surroundings. The spots on extreme horizontal branch stars are also significantly larger than sunspots, covering up to a quarter of the star's surface. These spots are incredibly persistent, lasting for decades, while individual sunspots are temporary, lasting only a few days to months. As the hot stars rotate, the spots on the surface come and go, causing the visible changes in brightness.

Beyond the variations in brightness due to spots, the team also discovered a couple of extreme horizontal branch stars that showed superflares—sudden explosions of energy and another signpost of the presence of a [magnetic field](#). "They are similar to the flares we see on our own Sun, but ten million times more energetic," says study co-author Henri Boffin, an astronomer at ESO's headquarters in Germany. "Such behavior was certainly not expected and highlights the importance of magnetic fields in explaining the properties of these stars."

After six decades of trying to understand extreme horizontal branch stars, astronomers now have a more complete picture of them. Moreover, this finding could help explain the origin of strong magnetic fields in many white dwarfs, objects that represent the final stage in the life of Sun-like stars and show similarities to extreme horizontal branch stars. "The bigger picture though," says team member, David Jones, a former ESO Fellow now at the Instituto de Astrofísica de Canarias, Spain, "is that changes in brightness of all hot stars—from young Sun-like stars to old extreme horizontal branch stars and long-dead white dwarfs—could all be connected. These objects can thus be understood as collectively suffering from magnetic spots on their surfaces."

More information: A plague of magnetic spots among the hot stars of globular clusters, *Nature Astronomy* (2020). [DOI: 10.1038/s41550-020-1113-4](https://doi.org/10.1038/s41550-020-1113-4) ,
www.nature.com/articles/s41550-020-1113-4

Provided by ESO

Citation: Hot stars are plagued by giant magnetic spots (2020, June 1) retrieved 16 August 2024 from <https://phys.org/news/2020-06-hot-stars-plagued-giant-magnetic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--