

New insights into the energy levels in quantum dots

June 25 2020



An electron inside a quantum dot is raised by a photon (green waveform) to a higher energy level. The result is a so-called exciton, an excited state consisting of two electrons and one hole. By emitting a photon (green waveform), the system returns to the ground state (green path). In rare cases, a radiative Auger process takes place (red arrow): an electron stays in the excited state, while a photon of lower energy (red waveform) is emitted. Credit: RUB, Arne Ludwig



Researchers from Basel, Bochum, and Copenhagen have gained new insights into the energy states of quantum dots. They are semiconductor nanostructures and promising building blocks for quantum communication. With their experiments, the scientists confirmed certain energy transitions in quantum dots that had previously only been predicted theoretically: the so-called radiative Auger process. For their investigations, the researchers in Basel and Copenhagen used special samples that the team from the Chair of Applied Solid State Physics at Ruhr-Universität Bochum had produced. The researchers report their results in the journal *Nature Nanotechnology*, published online on 15 June 2020.

Lock up charge carriers

In order to create a quantum dot, the Bochum researchers use selforganizing processes in crystal growth. In the process, they produce billions of nanometer-sized crystals of, for example, indium arsenide. In these they can trap charge carriers, such as a single electron. This construct is interesting for <u>quantum communication</u> because information can be encoded with the help of charge carrier spins. For this coding, it is necessary to be able to manipulate and read the spin from the outside. During readout, <u>quantum information</u> can be imprinted into the polarization of a photon, for example. This then carries the information further at the speed of light and can be used for quantum information transfer.

This is why scientists are interested, for example, in what exactly happens in the quantum dot when energy is irradiated from outside onto the <u>artificial atom</u>.



Special energy transitions demonstrated

Atoms consist of a positively charged core which is surrounded by one or more negatively charged electrons. When one electron in the atom has a high energy, it can reduce its energy by two well-known processes: in the first process the energy is released in the form of a single quantum of light (a photon) and the other electrons are unaffected. A second possibility is an Auger process, where the <u>high energy</u> electron gives all its energy to other electrons in the atom. This effect was discovered in 1922 by Lise Meitner and Pierre Victor Auger.



Schematic representation of a charged exciton, i.e. an excited state consisting of



two electrons and one hole within a quantum dot. Credit: Arne Ludwig

About a decade later, a third possibility has been theoretically described by the physicist Felix Bloch: in the so-called radiative Auger process, the excited electron reduces its energy by transferring it to both, a light quantum and another electron in the atom. A semiconductor quantum dot resembles an atom in many aspects. However, for quantum dots, the radiative Auger process had only been theoretically predicted so far. Now, the experimental observation has been achieved by researchers from Basel. Together with their colleagues from Bochum and Copenhagen, the Basel-based researchers Dr. Matthias Löbl and Professor Richard Warburton have observed the radiative Auger process in the limit of just a single photon and one Auger electron. For the first time, the researchers demonstrated the connection between the radiative Auger process and quantum optics. They show that quantum optics measurements with the radiative Auger emission can be used as a tool for investigating the dynamics of the single electron.

Applications of quantum dots

Using the radiative Auger effect, scientists can also precisely determine the structure of the quantum mechanical energy levels available to a <u>single electron</u> in the quantum dot. Until now, this was only possible indirectly via calculations in combination with optical methods. Now a direct proof has been achieved. This helps to better understand the quantum mechanical system.

In order to find ideal quantum dots for different applications, questions such as the following have to be answered: how much time does an electron remain in the energetically excited state? What <u>energy</u> levels form a quantum dot? And how can this be influenced by means of



manufacturing processes?

Different quantum dots in stable environments

The group observed the effect not only in quantum dots in indium arsenide semiconductors. The Bochum team of Dr. Julian Ritzmann, Dr. Arne Ludwig and Professor Andreas Wieck also succeeded in producing a quantum dot from the semiconductor gallium arsenide. In both material systems, the team from Bochum has achieved very stable surroundings of the quantum dot, which has been decisive for the radiative Auger process. For many years now, the group at Ruhr-Universität Bochum has been working on the optimal conditions for stable <u>quantum dots</u>.

More information: Matthias C. Löbl et al. Radiative Auger process in the single-photon limit, *Nature Nanotechnology* (2020). <u>DOI:</u> <u>10.1038/s41565-020-0697-2</u>

Provided by Ruhr-Universitaet-Bochum

Citation: New insights into the energy levels in quantum dots (2020, June 25) retrieved 27 June 2024 from <u>https://phys.org/news/2020-06-insights-energy-quantum-dots.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.