

Predator snails fend off the invasion of barnacles in Japan

June 3 2020, by Julius A. Ellrich and Takefumi Yorisue



Credit: Kevin B from Pexels

Introduced species can have tremendous ecological effects and [can become a major threat to ecosystems](#). Invasion biology tries to understand the mechanisms that [can limit and control the impacts of](#)

[introduced species.](#)

Through global shipping routes, barnacles have been [introduced to coastlines worldwide](#). An example is the barnacle *Balanus glandula*, native to the North American Pacific coast and [transported by ships to the Pacific coast of Honshu, Japan](#), where it has [replaced several native barnacle species](#). From Honshu, *B. glandula* spread to eastern Hokkaido (northern Japan), [where it occurs in relatively low densities compared to the native barnacle *Chthamalus dalli*](#).

Furthermore, *B. glandula* has reached as far as the Atlantic coasts of [Argentina](#) and [South Africa](#), with [negative effects](#) on native barnacle populations and other seafloor dwellers.

Biotic resistance is the ability of native species to limit the spread of invasive species. Working as marine ecologists at [Akkeshi Marine Station](#), on the Pacific coast of Hokkaido (Takefumi Yorisue, Kyosuke Momota), and the [Marine Ecology Lab](#), Saint Francis Xavier University in Nova Scotia (Julius A. Ellrich), we established an international collaboration to investigate the low *B. glandula* densities in Hokkaido rocky intertidal habitats.




Introduced barnacles (*Balanus glandula*, white arrow) and native barnacles (*Chthamalus dalli*, brown arrow). Credit: Takefumi Yorisue

Predatory and herbivorous snails

In Hokkaido, the native predatory snails *Nucella lima*—more commonly known as dogwhelks—and herbivorous snails, *Lottia cassis*, known as limpets, are common. Dogwhelks have a particular appetite for barnacles: [previous research in Atlantic Canada revealed that dogwhelks have nonconsumptive effects \(NCEs\)—behaviours that do not include consuming the barnacles—that can limit the appearance of new barnacles](#). These NCEs are triggered by dogwhelk chemical cues that limit barnacle cyprid larvae settlement. Cyprids swim away from dogwhelk cues [to reduce the risk of being consumed by dogwhelks](#).

Research has shown that as limpets graze, [they can accidentally ingest or disturb cyprids](#). These actions, known as disturbance effects (DEs), also limit [the appearance of new barnacles](#).

We hypothesised that the native dogwhelks and limpets would contribute to biotic resistance against *B. glandula*. To test our hypothesis, we conducted a series of experiments in Akkeshi Bay, in Hokkaido. We examined [dogwhelk feeding preferences for barnacles in the laboratory](#) and [dogwhelk NCEs and limpet DEs on barnacle population growth in the field](#).



The native dogwhelk *Nucella lima* (left) and the native limpet *Lottia cassis* (right). Credit: Takefumi Yorisue


Native dogwhelks and limpets

First, we compared dogwhelk feeding preferences for introduced and native barnacles by providing dogwhelks with similar-sized *B. glandula* and *C. dalli* individuals. We found that the dogwhelks preferred the introduced *B. glandula* over the local *C. dalli*. Next, we compared dogwhelk feeding preferences for large and small *B. glandula* individuals. We found that dogwhelks preferred large *B. glandula* individuals.

These preferences are likely driven by prey profitability: previous

research with dogwhelks on the North American Pacific coast indicated that [dogwhelks that fed on *B. glandula* grew faster than those who fed on *C. dalli*](#), and [dogwhelks fed with large *B. glandula* individuals grow the fastest](#).

Our findings suggest that dogwhelk predation limits the *B. glandula* spread in Hokkaido, [especially as large *B. glandula* individuals produce more offspring than small *B. glandula* individuals](#).



An illustration showing the effects of the presence of local dogwhelk on introduced barnacle populations. Credit: Julius A. Ellrich, Takefumi Yorisue and Kyosuke Momota

Population control measures

We examined the effect of dogwhelk NCEs on *B. glandula* settlement and population growth. For that, we installed cages with mesh compartments in the harbour of Akkeshi Marine Station. The cages prevented the dogwhelks from feeding on the barnacles, and we found that [the presence of dogwhelks limited the growth of *B. glandula* populations by 56 percent](#).

We conducted a similar experiment using limpets in a cage with the barnacles. We found that [feeding limpets limited the growth of *B. glandula* populations by 81 percent](#).

Employing native dogwhelks and limpets

Our results highlight that in the case of [barnacles](#) in Japan, predation and disturbance by native species are important ecological processes that can contribute to biotic resistance against introduced species. It is essential to understand such complex processes in order to protect native species that help to maintain biotic resistance. Without biotic resistance, it is more likely that introduced species—such as *B. glandula*—can spread and have negative effects on native [species](#) and communities.

Understanding biotic resistance can support local organisms in protecting their ecosystems against biological invasions, eliminating the need for more extreme measures.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Predator snails fend off the invasion of barnacles in Japan (2020, June 3) retrieved 11 May 2024 from <https://phys.org/news/2020-06-predator-snails-fend-invasion-barnacles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.