

Probing dark matter with the Higgs boson

June 15 2020



Figure 1: Mass of the leading two jets (x-axis) in the search region with all background processes stacked and compared to data. A hypothetical Higgs boson signal decaying to invisible final states is shown in red. Credit: ATLAS Collaboration/CERN

Visible matter—everything from pollen to stars and galaxies—accounts for roughly 15% of the total mass of the universe. The remaining 85% is



made of something entirely different from things we can touch and see: dark matter. Despite overwhelming evidence from the observation of gravitational effects, the nature of dark matter and its composition remain unknown.

How can physicists study dark matter beyond gravitational effects if it is practically invisible? Researchers are pursuing three approaches:

- indirect detection with astronomical observatories searching for the decay products of dark-matter annihilation in galactic centres
- direct detection with highly sensitive low-background experiments looking for dark matter scattering off nuclei
- creating dark matter in the controlled laboratory environment of the Large Hadron Collider (LHC) at CERN.

Although successful at describing <u>elementary particles</u> and their interactions at low energies, the Standard Model of particle physics does not include a viable dark-matter particle. The only possible candidates, neutrinos, do not have the right properties to explain the observed dark matter. To remedy this problem, a simple theoretical extension of the Standard Model posits that existing particles, such as the Higgs <u>boson</u>, act as a "portal" between known particles and dark-matter particles. Since the Higgs boson couples to mass, massive dark-matter particles should interact with it. The Higgs boson still has large uncertainties associated with the strength of its interaction with Standard Model particles; up to 30% of the Higgs-boson decays can potentially be invisible, according to the latest <u>ATLAS combined Higgs-boson</u> <u>measurements</u>.

Could some of the Higgs bosons decay into dark matter? As dark matter does not interact directly with the ATLAS detector, physicists look for signs of "invisible particles," inferred through momentum conservation of the proton–proton collision products. According to the Standard



Model, the fraction of Higgs bosons decaying to an invisible final state (four neutrinos!) accounts for just 0.1% and is thus negligible. Should such events be observed, it would be a direct indication of new physics and potential evidence of Higgs bosons decaying into dark-matter particles.



Figure 2: Upper limit on the WIMP-nucleon cross section at 90% confidence level derived in this analysis compared to direct detection experiments. Credit: ATLAS Collaboration/CERN

At the LHC, the most sensitive channel to search for direct decays of the Higgs boson to invisible particles is via the so-called vector boson fusion (VBF) production of the Higgs boson. VBF Higgs-boson production results in two sprays of particles (called "jets") that point in a more



forward direction in the ATLAS detector. This, combined with a large missing momentum in perpendicular direction ("transverse") to the beam axis from the invisible dark-matter particles, creates a unique signature that ATLAS physicists can search for.

The ATLAS Collaboration has studied the full LHC Run 2 dataset, collected by the detector in 2015–2018, to search for Higgs-boson decays to dark-matter particles in VBF events. No significant excess of events over the expected background from known Standard Model processes was found in the analysis. ATLAS derived, at a 95% confidence level, an exclusion bound of the Higgs-boson decay to invisible particles of 13%. This analysis included roughly 75% more data than the previous ATLAS search, and the team implemented several improvements including:

- Faster filtering algorithms to generate more simulated collisions with equivalent computing power. Lack of simulated events was the leading uncertainty in the first 13 TeV version of this analysis.
- Optimised collision selection to accept ~50% more Higgs-boson events on the same dataset.
- Refined event categorisation to result in a higher signal-tobackground ratio in the search regions. This can be seen in Figure 1 as the red curve in the lower panel increases with higher invariant mass of the two leading jets (m_{ii}).
- Improved acceptance for collisions enriched in background processes, allowing the analysts to improve the background-process modelling.

This observed exclusion is consistent with no signs of the Higgs boson decaying to dark matter. The new results advance the search for weakly interacting massive particles (WIMPs), a popular candidate for dark matter. ATLAS set additional exclusion limits for lower WIMP masses,



which are compared to other direct-detection experiments in Figure 2. These limits are competitive with the best direct-detection experiments for WIMP masses up to half of the Higgs-boson mass, assuming the Higgs boson interacts directly with dark matter.

This new analysis places the strongest existing limits on the Higgs boson decaying to invisible particles to date. As the search goes on, physicists will continue to increase the sensitivity to this fundamental probe of <u>dark</u> <u>matter</u>.

More information: Search for invisible Higgs boson decays with vector boson fusion signatures with the ATLAS detector using an integrated luminosity of 139 fb–1: <u>atlas.web.cern.ch/Atlas/GROUPS ...</u> <u>ATLAS-CONF-2020-008/</u>

Provided by ATLAS Experiment

Citation: Probing dark matter with the Higgs boson (2020, June 15) retrieved 9 May 2024 from <u>https://phys.org/news/2020-06-probing-dark-higgs-boson.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.