

A sharper view of flood risk

June 8 2020



More flexible statistical models could help improve the prediction of extreme rainfall events. Credit: Philip Scalia / Alamy Stock Photo

By generalizing a classical statistical model and adapting it for use in analyzing the extremes of rainfall in large datasets, researchers, including KAUST's Raphaël Huser, have devised a more efficient and flexible analytical tool that promises to improve the prediction of flood risk and other extreme weather phenomena.

Rare extreme weather events, such as floods, extreme winds, high temperatures and drought, can be devastating, but predicting the



frequency and severity of such conditions remains one of the key challenges in statistical science. Even large, long-term data sets over extensive areas may include very few extreme events, making it exceptionally difficult to predict future events with accuracy.

"There are classically two ways to model extreme events, the 'block maximum' approach, where we look at the largest events in blocks of time and the 'threshold exceedance' approach, which selects the top few percent of events across the entire timeframe of the dataset," explains Huser, who undertook the work in collaboration with U.S.-based colleagues Gregory Bopp and Benjamin Shaby. "Previous work has developed new tools to apply the threshold exceedance approach; in this study we generalized a classical block maximum model for application to extreme precipitation."

The block maximum approach has a long tradition in the statistics of extremes, but it has a high computational cost that limits its application to the large-scale datasets now routinely acquired in weather prediction. This approach is also unable to capture the observed weakening of the dependence between nearby conditions as events become more extreme.

The team's approach addresses both of these shortcomings by adapting a relatively inflexible, but computationally efficient, max-stable model using Bayesian inference, which is a statistical estimation approach that provides a natural way of incorporating expert opinion and accounting for various sources of variability.

"Our Bayesian model has a lot of parameters and hidden random effects, which need to be jointly estimated," says Huser. "Beyond the computational challenge, simply developing the model itself and deriving its theoretical properties was a major challenge. There is a reason why classical max-stable models have been used extensively for a long time—it is not straightforward to generalize them and come up with



more realistic and flexible models."

The model was able to capture the observed patterns in extreme precipitation events occurring along the coast and mountain range borders in northeast America, demonstrating its potential for predicting flood risk.

"Our <u>model</u> could also easily be adapted to other types of environmental datasets, such as wind and temperature, giving it very wide applicability," notes Huser.

More information: Gregory P. Bopp et al. A Hierarchical Max-Infinitely Divisible Spatial Model for Extreme Precipitation, *Journal of the American Statistical Association* (2020). DOI: <u>10.1080/01621459.2020.1750414</u>

Provided by King Abdullah University of Science and Technology

Citation: A sharper view of flood risk (2020, June 8) retrieved 16 June 2024 from <u>https://phys.org/news/2020-06-sharper-view.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.