

# Burrowing crabs reshaping salt marshes, with climate change to blame

July 14 2020

---



Credit: Christine Angelini

A new study reveals how climate change has enabled a voracious crab species to dramatically alter salt marsh ecosystems across the southeastern U.S.

The study, published in *Proceedings of the National Academy of Sciences*,

shows that soils beneath [salt marshes](#) from South Carolina to Florida have been softened by [higher sea levels](#) and increased tidal inundation. That softening has allowed the burrowing crab species *Sesarma reticulatum* to thrive, feeding on the cordgrass that holds the [marshes](#) together.

The clearing of grass by crabs has dramatically altered the flow of creeks that run through the marshes, the study found, and is altering the dynamics between predator and [prey species](#) in the marshes. In fact, the researchers say that *Sesarma*, which had previously been a minor player in southeastern salt marshes, can now be considered a keystone species, meaning it plays a dominant role in shaping the ecosystem.

"What we've found is an example of how [sea level rise](#) can activate a keystone species that's now dramatically remodeling these salt marshes," said Mark Bertness, a professor emeritus of ecology and evolutionary biology at Brown University and a coauthor of the research. "That's a big deal because sea level rise is a pervasive global phenomenon, and this is a largely unexpected consequence. We need to start thinking about how [global climate change](#) could activate new keystone species in other ecosystems."

Research on *Sesarma* crabs and their impact on salt marshes has a long history in Bertness's lab at Brown. In 2011, Bertness and his students discovered that *Sesarma*, voracious grazers of cordgrass roots and leaves, were behind sudden die-offs of marshes on Cape Cod. In that case, overfishing had suddenly pulled predator species like striped bass out of the water, giving the crabs free reign to decimate the marshes. One of the undergraduate co-authors on that earlier research was Christini Angelini, now an associate professor at the University of Florida and a senior author on this new paper.

*Sesarma* were known to inhabit southern marshes in Florida and the

Carolinas, but their populations hadn't boomed like those further north. One potential reason for that was differing soil substrates. While working several years ago as an undergraduate researcher in Bertness' lab, Sinead Crotty, now project director at Yale's Carbon Containment Lab, showed that ground hardness played a big role in where *Sesarma* are able to establish themselves. Her findings indicated that *Sesarma* had a much easier time building burrows and feeding on grass roots in the peaty New England soil compared to harder soil substrates often found in southern marshes.



The image shows a marsh creekhead cleared of its cordgrass by the voracious grazing of *Sesarma* crabs. The clearing of grasses increases the erosion rates of marshland Credit: Christine Angelini

But as sea levels continue to rise due to climate change, Crotty, Angelini and Bertness wondered if softening soils might be giving *Sesarma* more of a foothold in the South. Looking at aerial photos from nine locations across South Carolina and Florida, they found that the number of marsh

creeks with evidence of *Sesarma* grazing increased by up to 240% from the late 1990s to the late 2010s. Meanwhile, surveys of sea level rise show that the ground in these areas is tidally submerged up to an hour longer per day now compared to the late 1990s.

"You've got the sea level rising, which softens the substrate that these crabs usually can't burrow in," Bertness said. "Now that it's softer you've got an ideal habitat to support these huge communal *Sesarma* burrows."

This new *Sesarma* activity is reshaping marshes, the researchers found. Elimination of grasses has increased the rate at which creeks form in the marshes, and increases the drainage density of marsh creeks by up to 35%.

*Sesarma* activity is also influencing interactions between predators and prey in the creeks. Clearing of grasses provides predators increased access to shellfish and other prey species. The research found that populations of mussels were dramatically lower in *Sesarma*-grazed creeks compared to creeks that weren't grazed.

"As they drown, southeastern U.S. marshes are fracturing from grasslands to patches of marsh, with depleted populations of mussels, snails and other invertebrates," Angelini said. "These dynamics reveal how quickly marshes may disappear with accelerating sea level rise and how long they will remain foraging grounds for commercially, recreationally and ecologically important species."

The fact that *Sesarma* is now altering the geomorphology of the marshes, as well as the ecological interactions between other species, is evidence that it now qualifies as a keystone species in southern marshes. This is the first example, the researchers say, of activation of new keystone species as the result of anthropogenic climate change.

"This is going to be something for the textbooks," Bertness said. "This is an underappreciated way in which [climate change](#) alters ecosystems."

**More information:** Sinéad M. Crotty et al, Sea-level rise and the emergence of a keystone grazer alter the geomorphic evolution and ecology of southeast US salt marshes, *Proceedings of the National Academy of Sciences* (2020). [DOI: 10.1073/pnas.1917869117](https://doi.org/10.1073/pnas.1917869117)

Provided by Brown University

Citation: Burrowing crabs reshaping salt marshes, with climate change to blame (2020, July 14) retrieved 25 April 2024 from

<https://phys.org/news/2020-07-burrowing-crabs-reshaping-salt-marshes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.