


Cell death in porpoises caused by environmental pollutants

July 20 2020



Cell death and risk assessment of finless porpoise fibroblasts by exposure to environmental pollutants. Credit: American Chemical Society

A recent study just published in *Environmental Science & Technology* identified the toxicological risks of environmental pollutants to finless

porpoises (*Neophocaena asiaeorientalis*). Manmade chemicals synthesized for human activities threaten the health of marine mammals. These chemicals, including persistent organic pollutants (POPs), have long been known to accumulate at high levels in many dolphin species. The POPs levels of finless porpoises inhabiting the Seto Inland Sea are higher than those of other cetacean species distributed in the waters near Japan, and the effects of toxicity have been a concern. Nevertheless, ecotoxicological studies of wild dolphins are difficult due to legal and ethical considerations, and information is lacking. Researchers in the Center for Marine Environmental Studies (CMES), Ehime University, together with collaborators, have successfully isolated the fibroblast cells from a finless porpoise stranded in the Seto Inland Sea, Japan revealing the toxicological risk of pollutants of concern in the local population.

Cell culture and exposure to pollutants

Fibroblasts of a finless porpoise were collected from a stranded individual. Seventeen chemicals including dioxin (2,3,7,8-tetrachlorodibenzo-p-dioxin, TCDD), industrial chemicals (polychlorinated biphenyls, PCBs), metabolites of PCBs (hydroxylated PCBs, OH-PCBs), flame retardants (polybrominated biphenyls, PBDEs), insecticides (dichlorodiphenyltrichloroethane and their metabolites, DDTs), and methylmercury were tested for their cellular toxicities.

Effects of pollutants on fibroblasts

Most pollutants induced cell death at higher concentrations, and dioxin-like compounds (TCDD and dioxin-like PCBs) were more toxic than the other chemicals tested. Toxic potencies of OH-PCBs and its precursor PCBs were different for each endpoint, and these compounds may contribute to cell damage by different mechanisms. Dose-dependent cell damage was also observed with DDTs, which accumulated in relatively

high concentrations in many whale species. Among DDTs, *p,p'*-DDT was the most potent for the cytotoxicity, whereas *p,p'*-DDE notably affected the cell viability. Methylmercury also induced cellular necrosis at the highest test concentration (100 μ M).

Risk assessment at the population level

To assess the risk of the porpoise population inhabiting the Seto Inland Sea, the research group estimated the EARs (exposure-activity ratios). EAR is the emerging concept of finding high-risk [chemical](#) substances by comparing the concentrations at which cytotoxicity was observed with the concentration of the chemicals in animal bodies. Collectively, PCBs and DDTs were shown to be at high risks and could cause cytotoxicity, apoptosis, and reduced cell viability in the porpoise population in the Seto Inland Sea.

This study successfully evaluated the risks of [environmental pollutants](#) using fibroblasts isolated from a dead [porpoise](#). There is an urgent need to better and comprehensively understand the risks of pollutants not only in this species but also in other marine mammals, and it is important to implement measures to reduce the load of high-risk pollutants in the marine environment.

More information: Mari Ochiai et al. In Vitro Cytotoxicity and Risk Assessments of Environmental Pollutants Using Fibroblasts of a Stranded Finless Porpoise (*Neophocaena asiaeorientalis*), *Environmental Science & Technology* (2020). [DOI: 10.1021/acs.est.9b07471](https://doi.org/10.1021/acs.est.9b07471)

Provided by ResearchSEA

Citation: Cell death in porpoises caused by environmental pollutants (2020, July 20) retrieved 26 June 2024 from

<https://phys.org/news/2020-07-cell-death-porpoises-environmental-pollutants.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.