

Where is water distributed during a drought?

July 17 2020



Dried out stream bed in Demnitz, Germany. Credit: Lukas Kleine

In low precipitation periods, where and how is the limited available water distributed, and what possibilities are there for improving retention in the soil and the landscape? Dörthe Tetzlaff and her team from the Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB) have discovered that vegetation has a major influence on this. The

researchers are investigating the storage, distribution and quality of water in the landscape. Using the example of the drought-sensitive Demnitzer Mühlenfließ in Brandenburg, a sub-catchment area of the Spree, they quantified visible and invisible water flows during and shortly after the drought of 2018.

The annual rainfall in Brandenburg is 560 liters per square meter. This makes Brandenburg a region with the lowest rainfall in Germany. In 2018 there were 390 liters of water per square meter, which is about 40 percent less precipitation than usual.

Even under "normal" climatic conditions, about 90 percent of the precipitation is released back into the atmosphere and does not flow into groundwater or rivers. Groundwater levels in the area today show that the decreased water levels due precipitation deficits from 2018 could not be returned to normal conditions between the growing seasons.

Land use critical for water resource management

Dörthe Tetzlaff is a researcher at the IGB and Professor for Ecohydrology at the Humboldt University of Berlin. She and her team investigated how the process of evaporation and groundwater recharge differ under different soils and land uses.

"Due to the current climate crisis with increasing droughts, we need to know how much water different plants use. As researchers, we ask ourselves: Can we apply sustainable land use to control water consumption and make entire landscapes more resilient towards climate extremes? These findings are the basis for meeting the demands for food production and water supply," says Dörthe Tetzlaff explaining her motivation for her research topic.

Forest soil drier than grassland

In the Demnitzer Mühlenfließ, the team investigated two sites with land uses typical for the region: a mixed forest site with [sandy soils](#) and a deep rooting zone; and grassland site with loamier soils and a shallower rooting zone. The forest [soil](#) was much drier, which is due to the characteristics of soil and plants.

For example, during the drought, the uppermost meter of the sandy soil in the forest contained only 37 liters per m^2 , and under grassland there were as much as 146 liters of water per m^2 . The treetops/leaf canopy of the forest shielded part of the rain that evaporated directly from the leaves and never reached the ground. Moreover, the sandy forest soil caused faster water flow through the soil and decreased water storage. Rainfall penetrated deeper into the soil, but was reabsorbed by the trees during the growing season before reaching the groundwater.

Under the grassland, the water continuously recharged the groundwater. The soil could store more water. As the plants only took water from the upper soil, this led to "older" soil water.

"We were able to show how poorly the landscapes in Brandenburg store precipitation, limiting drought resistance. The type of forest we investigated is typical for the Northern European Plain. It was sad to see that even a natural mixed forest is suffering greatly from drought. For economically used forests that are dominated by conifers, the situation is even worse. In fact, the conifer mortality in Brandenburg is now obvious," says Lukas Kleine, doctoral student in Tetzlaff's team.

"Planting water"—How agriculture can take advantage of the research results

The researchers work together with the agricultural and forestry sectors to bring their research results into practice. One of their most important partners is Benedikt Bösel, owner of Gut & Bösel. The agricultural holding Gut & Bösel tests and develops multifunctional land use concepts of regenerative agriculture and forestry and confirms the observations of the IGB researchers: "The regeneration of our soils and soil health is the greatest and most important task of our generation. For this we need innovative solutions in agriculture and forestry to fight the causes of our problems instead of just curing the symptoms. Only in this way can we act in accordance with the complexity of the ecosystems. We are trying to develop these solutions based on the findings of Prof. Tetzlaff's team, among others".

"We see that after the further dry seasons in 2019 and so far in 2020, the groundwater levels will continue to fall. The vegetation has still not been able to recover due to the low rainfall in the winter months. Unfortunately we are far from 'normal' conditions. In order to improve the resistance of Brandenburg's ecosystems to droughts and other climate changes, measures must be implemented that promote groundwater recharge and create soils that can store more water. Our results underline the central role of vegetation in the development of such strategies," says Dörthe Tetzlaff.

Water in the landscape: Blue and green water

Researchers distinguish between the so-called blue water, which fills lakes, rivers and aquifers and is immediately available for water supply; and the green water, which is directly influenced by vegetation and is returned to the atmosphere through evaporation and transpiration after being absorbed by plants. Dörthe Tetzlaff and her team are investigating the interactions between blue and green water fluxes. They are analyzing what happens in the critical zone and what influence vegetation has on the overall water regime.

The critical zone—the thin, dynamic and life-supporting skin of the earth

The layer of earth that extends between the canopy, the soil and the groundwater is called the critical zone. For a long time it was a 'black box'; in particular, the role of plants in the distribution of water has been neglected, as science has focused on the blue water fluxes.'

In this study, the researchers investigated the water fluxes in the critical zone with stable isotopes in the water. Stable isotopes in water can be used as "markers" to determine flow paths, age and origin of water. For a comprehensive understanding, it is not only the absolute amounts of the water flows in the landscape that are important, but also how long the water is stored on site and which flow paths it takes. When this information is integrated with data on vegetation dynamics, tracer-based modeling can reveal some of the most important processes in the ecohydrological system, such as where and at what rate plants draw [water](#) from the soil.

More information: Lukas Kleine et al. Using isotopes to understand evaporation, moisture stress and re-wetting in catchment forest and grassland soils of the summer drought of 2018, (2020). [DOI: 10.5194/hess-2020-81](https://doi.org/10.5194/hess-2020-81)

Aaron Smith et al. Isotope-aided modelling of ecohydrologic fluxes and water ages under mixed land use in Central Europe: The 2018 drought and its recovery, *Hydrological Processes* (2020). [DOI: 10.1002/hyp.13838](https://doi.org/10.1002/hyp.13838)

Provided by Forschungsverbund Berlin e.V. (FVB)

Citation: Where is water distributed during a drought? (2020, July 17) retrieved 23 May 2024 from <https://phys.org/news/2020-07-drought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.