

The restoration of forests with active rapid 'Ohi'a death infections may be possible

July 9 2020



'Ohi'a (*Metrosideros polymorpha*) seedling in an unweeded plot. Credit: Stephanie Yelenik, Pacific Island Ecosystem Research Center. Public domain.

For the first time, researchers have shown that native 'ohi'a seedlings can

survive for at least a year in areas that have active mortality from Rapid 'ohi'a Death, or ROD, a fungal disease that is devastating to this dominant and culturally important tree in Hawaiian forests. This information can be useful to land managers and homeowners as they prioritize conservation actions.

The study, published recently in *Restoration Ecology*, was authored by scientists from the U.S. Geological Survey and Hawai'i Cooperative Studies Unit at University of Hawai'i Hilo.

"'Ohi'a is a keystone species in Hawaiian forests, and ROD has the potential to cause major ecosystem disturbances that will negatively impact water supply, cultural traditions, natural resources and quality of life," said USGS Director Jim Reilly. "This [innovative research](#) provides a glimmer of hope for native 'ohi'a tree restoration in Hawaii by indicating that successful planting of 'ohi'a could be possible in ROD-affected forests if the native species' seedlings are protected."

The study also highlights specific best practices for maximizing seedling survival, noted Stephanie Yelenik, an ecologist with the USGS and lead author of the study.

"We found that 'ohi'a seedlings planted into a [forest](#) heavily affected by ROD have a high probability of survival for the first year," said Yelenik. "While that one-year survival of seedlings is great news, this species lives centuries and there's currently no treatment once the tree becomes infected. Because 'ohi'a grow slowly, a dead tree is a gap in the canopy for a long time, and one of Hawai'i's many quick-growing invaders can take over the gaps caused by dead trees."

Still, she added, "these results provide the first, but still early, evidence that planting 'ohi'a may be used as a restoration tool in forests that are threatened by ROD.

ROD is a newly emerging disease caused by fungal pathogens. Since the disease was first discovered, ROD has killed at least 1 million 'ohi'a—a tree that is foundational to the Hawaiian landscape and culture. Its loss, study authors emphasized, is detrimental to habitat for endangered birds and plants, ecosystem processes and the cultural heritage of Hawaiians. There remains no treatment for infected trees, so slowing the spread of ROD remains important to protect native Hawaiian forests.

"The 'ohi'a tree is the dominant tree in Hawaiian forests, the first to colonize new lava flows and often forms the bulk of what makes up the forests," said Yelenik. "While planting 'ohi'a may not be necessary in forests where 'ohi'a seedlings naturally occur, we thought planting might be a good tool for managers who want to maintain a native 'ohi'a canopy in forests dominated by invasive plants in the understory. We wanted to test that idea."

To test how 'ohi'a seedlings would fare in areas where ROD is prevalent, researchers planted seedlings directly underneath adult 'ohi'a that had tested positive for *Ceratocystis huliohia* or *lukuohia*, the fungal pathogens that cause ROD. They monitored survival in plots that were fenced to keep animals out and removed seedlings of invasive trees and shrubs; other seedlings were planted in areas with no fencing or weeding. Seedlings that died were brought back to the lab for testing. In the lab researchers were able to test all the seedlings for the DNA of *Ceratocystis* and tested for evidence of viable fungal spores.

The results were clear: none of the dead seedlings tested positive for *Ceratocystis*, indicating that 'ohi'a can survive in ROD-affected forests if protected from wild pigs and goats and invasive trees and shrubs. In addition, seedlings were six times more likely to die in plots where weeds were allowed to grow than in areas where weeds were cleared around the 'ohi'a seedlings. In terms of prioritizing management, this means that controlling [invasive plants](#) and animals has a greater impact

on survival in the tree's first year than does exposure to ROD.

"The current study provides clues about the epidemiology of the disease and the dynamics of ROD pathogens across the landscape," said Lisa Keith, research plant pathologist for the USDA Agricultural Research Service. "This information can help to slow the spread of the disease through effective management strategies. While surrounding disease pressure may be high, results suggest that ohia seedlings have the capacity to thrive in areas devastated by ROD."

Results of the study indicated that 'ohi'a seedlings are not infected with ROD through the soil. "Those results are encouraging for two reasons," said J.B. Friday, extension forester with the University of Hawai'i Cooperative Extension Unit. "First, it means that even in forests with invasive [trees](#) and shrubs, 'ohi'a may possibly be re-established. And second, it means that in our high-elevation, pristine native forests, natural 'ohi'a regeneration could be possible, even in forests hit by ROD, if those areas are protected."

Yelenik cautioned that longer-term studies of 'ohi'a seedling survival in ROD-affected forests are still needed, but these early results demonstrate that active planting could successfully help maintain native 'ohi'a forests. Survival of 'ohi'a [seedlings](#) in ROD-affected forests is good news, but the author noted that "protecting 'ohi'a from infection remains the primary tool in the fight against ROD."

More information: Stephanie Yelenik et al, Successful restoration of *Metrosideros polymorpha* ('Ōhi'a) is possible in forest sites with active Rapid 'Ōhi'a Death infections, *Restoration Ecology* (2020). [DOI: 10.1111/rec.13197](https://doi.org/10.1111/rec.13197)

Provided by United States Geological Survey

Citation: The restoration of forests with active rapid 'Ohi'a death infections may be possible (2020, July 9) retrieved 3 May 2024 from <https://phys.org/news/2020-07-forests-rapid-ohia-death-infections.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.