

Light pollution gives invasive cane toads a belly full of grub

July 1 2020



Relationship between light pollution and invasive toads' foraging behavior. Artificial light provides food resources such as insects for invasive toads, and the amount is influenced by lunar phases. Credit: Hirotaka Komine/ TUAT

Light pollution is increasingly being recognized as yet another of humanity's many impacts on the rest of the natural world. Artificial light at night, or 'ALAN' to use the researchers' name for the phenomenon,



can alter foraging behavior, migration patterns, mortality rates and even the very physical structure of animals.

But there has been little research into the effect of ALAN on invasive species—species that are not indigenous, or native, to an area—which can sometimes cause significant economic and ecological damage.

However, as anyone who has a porch <u>light</u> can see, nocturnal insects such as moths, flies and beetles are attracted to this artificial light. These invertebrates are <u>food sources</u> for nocturnal predators such as bats, birds and frogs—including what is perhaps the most iconic invasive species of all time, the cane toad.

Cane toads feed on nocturnal insects, and Assistant Professor Hirotaka Komine with Tokyo University of Agriculture and Technology (TUAT) along with their colleagues from Australia's James Cook University wanted to find out whether the artificially large food source that ALAN makes available in a particular location had any impact on the cane toad.

They published their findings on April 16 in Scientific Reports.

They set up six outdoor enclosures, each with an artificial light attached, in different rural areas around Townsville, Queensland. They then placed <u>cane toads</u> that they had locally captured in the enclosures just before sunset. After the toads had feasted overnight on whatever invertebrate delectable treats had been attracted to the ALAN, the toads were euthanized and dissected to measure the contents of their bellies.

The researchers varied the amount of light, and controlled for the affects of rainfall, <u>wind speed</u>, and temperature, which also have an impact on the availability of sources of food. They found that when the lights were on, the percentage of flying insects that the toads consumed increased significantly.



But the toads did not eat as many flying bugs when the moon was bright. The same reduction in bug consumption happened in those areas that were closer to urban areas that already suffered from ample ambient ALAN. In both cases, the extra light diluted the impact of the lamps the researchers had set up.

"This tells us that in order to control the growth of this <u>invasive species</u>," said conservation biologist Hirotaka Komine. "More careful management of artificial light in rural areas such as that from highway lampposts or scattered buildings, could go a long way in reducing the food resources available."

The researchers added that light management approaches should also consider the <u>lunar cycle</u>, such as reducing light use where possible during dark lunar phases.

More information: Hirotaka Komine et al, Impacts of artificial light on food intake in invasive toads, *Scientific Reports* (2020). DOI: 10.1038/s41598-020-63503-9

Provided by Tokyo University of Agriculture and Technology

Citation: Light pollution gives invasive cane toads a belly full of grub (2020, July 1) retrieved 16 June 2024 from <u>https://phys.org/news/2020-07-pollution-invasive-cane-toads-belly.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.