


Study reveals composition of 'gel-like' substance discovered by Chang'e-4 rover on Moon's far side

July 21 2020



Impact melt breccia and surrounding context Credit: CNSA, CLEP, and AIR


The unusual dark greenish and glistening 'gel-like' substance in a crater on the far side of the moon has attracted widespread interest following its discovery by the Chang'e-4 rover in July 2019.

A research team led by Prof. Di Kaichang from the Aerospace Information Research Institute (AIR) of the Chinese Academy of Sciences and their collaborators analyzed the substance in detail by using multiple datasets from the rover's panoramic camera (Pancam), hazard avoidance camera (Hazcam), and the visible and near-infrared spectrometer (VNIS).


The researchers found that the unusual substance is actually an impact melt breccia, and the provenance of the rover measured surrounding regolith might originate from a differentiated melt pool or from a suite of igneous rocks. Their findings were published in *Earth and Planetary Science Letters*.

China's Chang'e-4 probe, including a lander and a rover, successfully touched down within the 185-km-wide Von Kármán crater inside the South Pole-Aitken (SPA) basin on January 3, 2019, making the first-ever [soft landing](#) on the lunar far side.

The gel-like substance was discovered in a crater during the eighth lunar day of the rover's mission. Detailed measurements on the breccia and surrounding regolith were conducted during the ninth lunar day.



Photos of lunar impact melt breccia samples Credit: NASA and AIR



Hazcam images show the rover's locations Credit: CNSA, CLEP, and AIR

The discovered breccia, measuring 52 by 16 centimeters, resembles the lunar impact melt breccia samples 15466 and 70019 returned by the Apollo missions. It was formed by impact-generated welding, cementing and agglutinating of lunar regolith and breccia.

Clods surrounding the breccia-hosting crater were crushed into regolith powders by the rover's wheels, indicating the regolith may be compacted slightly and becomes blocky and friable. Hapke model-based spectral unmixing results showed that plagioclase was abundant, and pyroxene and olivine had almost equal fractions, indicating the regolith was likely the weathering products of noritic rocks.

The regolith measured by Chang'e-4 rover was actually a mixture of multiple sources, with ejecta from Finsen crater being primary and possible contributions from Alder [crater](#). Finsen and Alder craters are on the margin of the proposed impact melt pool produced by the SPA basin-forming event. Therefore, the provenance of the [regolith](#) might originate from a differentiated melt pool or from a suite of igneous rocks.

More information: Sheng Gou et al, Impact melt breccia and surrounding regolith measured by Chang'e-4 rover, *Earth and Planetary Science Letters* (2020). [DOI: 10.1016/j.epsl.2020.116378](https://doi.org/10.1016/j.epsl.2020.116378)

Provided by Chinese Academy of Sciences

Citation: Study reveals composition of 'gel-like' substance discovered by Chang'e-4 rover on Moon's far side (2020, July 21) retrieved 23 April 2024 from <https://phys.org/news/2020-07-reveals-composition-gel-like-substance-change-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.